

# Euler.


Leonhard Euler (1707-1783)



- Successor of **Nicolaus Bernoulli** in St. Petersburg (1726-1727).
- Member of the newly founded St. Petersburg Academy of Sciences (1727).
- 1741-1766: Director of Mathematics, later inofficial head of the Berlin Academy.

# Euler diagrams.

*Lettres à une Princesse d'Allemagne (1768-72).*



"Every  $A$  is  $B$ ."

"No  $A$  is  $B$ ."

"Some (but only some)  $A$  is  $B$ ."


"Some (but only some)  $A$  is not  $B$ ."

(Diagrams with Existential import!)


# Gergonne (1).

Joseph Diaz Gergonne (1771-1859).

- Very active in the wars after the French revolution.
- Discoverer of the duality principle in geometry.
- *Essais de dialectique rationnelle* (1816-1817):



# Gergonne (2).



Any two non-empty extensions (“sets”)  $A$  and  $B$  are in exactly one of Gergonne’s five relations:

- $h$  *est hors de*
- $x$  *s’entre-croise avec*
- $l$  *est identique à*
- $c$  *est contenue dans*
- $\supset$  *contiens*

# Gergonne (3).

Syllogisms of the first figure:  $A \bullet_0 B, B \bullet_1 C : A \bullet_2 C$ .

	h	x	l	c	o
h		$\neg l, \neg o$	h	$\neg l, \neg o$	h
x	$\neg l, \neg c$		x	$\neg h, \neg l, \neg o$	$\neg l, \neg c$
l	h	x	l	c	o
c	h	$\neg l, \neg o$	c	c	
o	$\neg l, \neg c$	$\neg h, \neg l, \neg c$	o	$\neg h$	o

If  $AxB$  and  $BcC$ , then  $\neg AIC$  and  $\neg AoC$ .

# De Morgan.

Augustus de Morgan (1806-1871).



- Professor of Mathematics at UCL (1828).
- Corresponded with [Charles Babbage](#) (1791-1871) and [William Rowan Hamilton](#) (1805-1865).
- **1866.** First president of the [London Mathematical Society](#).
- $x = 43, x^2 = 1849. y = 45, y^2 = 2025.$


**De Morgan rules.**

$$\neg(\Phi \wedge \Psi) \equiv \neg\Phi \vee \neg\Psi$$

$$\neg(\Phi \vee \Psi) \equiv \neg\Phi \wedge \neg\Psi$$

# Boole (1).

George Boole (1815-1864).



- School teacher in Doncaster, Liverpool, Waddington (1831-1849).
  - Correspondence with [de Morgan](#).
  - Professor of Mathematics at Cork (1849).
- 
- Developed an [algebra of logic](#) based on the idea of taking the extensions of predicates as objects of the algebra.
  - 1 is the “universe of discourse”, 0 is the empty extension.

# Boole (2).

“All  $B$  are  $A$ ”  $b(1 - a) = 0.$

“No  $B$  is an  $A$ ”  $ba = 0.$

“Some  $B$  is an  $A$ ”  $ba \neq 0.$


“Some  $B$  is not an  $A$ ”  $b(1 - a) \neq 0.$

## Celarent.

- *We assume:*  $ba = 0$  and  $c(1 - b) = 0.$
- *We have to show:*  $ca = 0.$
- $ba = 0$  implies that  $cba = c0 = 0.$
- $ca = ca - 0 = ca - cba = a(c - bc) = a(c(1 - b)) = ac0 = 0.$

# Venn.

John Venn (1834-1923).



- Lecturer in Moral Science at Cambridge (1862).
- Area of interest: logic and probability theory.
- *Symbolic Logic* (1881).
- *The Principles of Empirical Logic* (1889).
- *Alumni Cantabrigienses*.

**Venn diagrams.**

# Boolean Algebras (1).

A structure  $\mathbf{B} = \langle B, 0, 1, +, \cdot, - \rangle$  is a **Boolean algebra** if

- $B$  is a set with  $0, 1 \in B$ .
- $+$  and  $\cdot$  are binary operations on  $B$  satisfying the commutative and associative laws.
- $-$  is a unary operation on  $B$ .
- $+$  distributes over  $\cdot$  and *vice versa*:  $x + (y \cdot z) = (x + y) \cdot (x + z)$  and  $x \cdot (y + z) = (x \cdot y) + (x \cdot z)$ .
- $x \cdot x = x + x = x$  (idempotence),  $- - x = x$ .
- $-(x \cdot y) = (-x) + (-y)$ ,  $-(x + y) = (-x) \cdot (-y)$  (de Morgan's laws).
- $x \cdot (-x) = 0$ ,  $x + (-x) = 1$ ,  $x \cdot 1 = x$ ,  $x + 0 = x$ ,  $x \cdot 0 = 0$ ,  $x + 1 = 1$ .
- $-1 = 0$ ,  $-0 = 1$ .

**Example.**  $B = \{0, 1\}$ .

	$\cdot$	0	1		$+$	0	1
0		0	0		0	0	1
1		0	1		1	1	1

# Boolean Algebras (2).

$X := \{\text{Platon, Aristotle, Speusippus, Themistokles}\}$


$\mathbf{Phil} := \{\text{Platon, Aristotle, Speusippus}\}$

$\mathbf{Rhet} := \{\text{Themistokles}\}$

$\mathbf{Acad} := \{\text{Platon, Speusippus}\}$

$\mathbf{Peri} := \{\text{Aristotle}\}$

$B := \{\emptyset, X, \mathbf{Phil}, \mathbf{Rhet}, \mathbf{Acad}, \mathbf{Peri}, \mathbf{Rhet} + \mathbf{Peri}, \mathbf{Rhet} + \mathbf{Acad}\}.$



# Boolean Algebras (3).

If  $X$  is a set, let  $\wp(X)$  be the **power set** of  $X$ , *i.e.*, the set of all subsets of  $X$ .

For  $A, B \in \wp(X)$ , we can define

- $A \cdot B := A \cap B,$
- $A + B := A \cup B,$
- $0 := \emptyset,$
- $1 := X,$
- $-A := X \setminus A.$

Then  $\langle \wp(X), 0, 1, +, \cdot, - \rangle$  is a Boolean algebra, denoted by  $\mathbf{Pow}(X)$ .

# Boolean Algebras (4).

Define the notion of isomorphism of Boolean algebras: Let  $\mathbf{B} = \langle B, 0, 1, +, \cdot, - \rangle$  and  $\mathbf{C} = \langle C, \perp, \top, \oplus, \otimes, \ominus \rangle$  be Boolean algebras. A function  $f : B \rightarrow C$  is a **Boolean isomorphism** if

- $f$  is a bijection,
- for all  $x, y \in B$ , we have  $f(x + y) = f(x) \oplus f(y)$ ,  
 $f(x \cdot y) = f(x) \otimes f(y)$ ,  $f(-x) = \ominus f(x)$ ,  $f(0) = \perp$ ,  
 $f(1) = \top$ .

**Stone Representation Theorem.** If  $\mathbf{B}$  is a Boolean algebra, then there is some set  $X$  such that  $\mathbf{B}$  is isomorphic to a subalgebra of  $\text{Pow}(X)$ .

# Circuits.

- + corresponds to having two switches in parallel: if either (or both) of the switches are **ON**, then the current can flow.
- · corresponds to having two switches in series: if either (or both) of the switches are **OFF**, then the current is blocked.

# Mathematics and real content.

*Mathematics getting more abstract...*

## **Imaginary numbers.**

Niccolo Tartaglia      Girolamo Cardano

(1499-1557)

(1501-1576)



# Mathematics and real content.

*Mathematics getting more abstract...*

## **Imaginary numbers.**

Nicolo Tartaglia    Girolamo Cardano  
(1499-1557)        (1501-1576)


Carl Friedrich Gauss (1777-1855)

## **Ideal elements in number theory.**

Richard Dedekind (1831-1916)



# The Delic problem (1).



If a cube has height, width and depth 1, then its volume is  $1 \times 1 \times 1 = 1^3 = 1$ .

If a cube has height, width and depth 2, then its volume is  $2 \times 2 \times 2 = 2^3 = 8$ .

In order to have volume 2, the height, width and depth of the cube must be  $\sqrt[3]{2}$ :

$$\sqrt[3]{2} \times \sqrt[3]{2} \times \sqrt[3]{2} = (\sqrt[3]{2})^3 = 2.$$


# The Delic problem (2).

**Question.** Given a compass and a ruler that has only integer values on it, can you give a geometric construction of  $\sqrt[3]{2}$ ?

**Example.** If  $x$  is a number that is constructible with ruler and compass, then  $\sqrt{x}$  is constructible.

*Proof.*

If  $x$  is the sum of two squares (i.e.,  $x = n^2 + m^2$ ), then this is easy by Pythagoras. In general:



# The Delic problem (3).

It is easy to see what a **positive solution** to the Delic problem would be. But a **negative solution** would require reasoning about all possible geometric constructions.

# Geometries (1).

- We call a structure  $\langle P, L, I \rangle$  a **plane geometry** if  $I \subseteq P \times L$  is a relation.
- We call the elements of  $P$  “**points**”, the elements of  $L$  “**lines**” and we read  $pI\ell$  as “ **$p$  lies on  $\ell$** ”.
- If  $\ell$  and  $\ell^*$  are lines, we say that  **$\ell$  and  $\ell^*$  are parallel** if there is no point  $p$  such that  $pI\ell$  and  $pI\ell^*$ .
- **Example.** If  $P = \mathbb{R}^2$ , then we call  $\ell \subseteq P$  a **line** if

$$\ell = \{ \langle x, y \rangle ; y = a \cdot x + b \}$$

for some  $a, b \in \mathbb{R}$ . Let  $\mathcal{L}$  be the set of lines. We write  $pI\ell$  if  $p \in \ell$ . Then  $\langle P, \mathcal{L}, I \rangle$  is a plane geometry.

# Geometries (2).

- (A1) For every  $p \neq q \in P$  there is exactly one  $\ell \in L$  such that  $pI\ell$  and  $qI\ell$ .
- (A2) For every  $\ell \neq \ell^* \in L$ , either  $\ell$  and  $\ell^*$  are parallel, or there is exactly one  $p \in P$  such that  $pI\ell$  and  $pI\ell^*$ .
- (N) For every  $p \in P$  there is an  $\ell \in L$  such that  $p$  doesn't lie on  $\ell$  and for every  $\ell \in L$  there is an  $p \in P$  such that  $p$  doesn't lie on  $\ell$ .
- (P2) For every  $\ell \neq \ell^* \in L$ , there is exactly one  $p \in P$  such that  $pI\ell$  and  $pI\ell^*$ .

A plane geometry that satisfies (A1), (A2) and (N) is called a **plane**. A plane geometry that satisfies (A1), (P2) and (N) is called a **projective plane**.

# Geometries (3).

- (A1) For every  $p \neq q \in P$  there is exactly one  $\ell \in L$  such that  $pI\ell$  and  $qI\ell$ .
- (A2) For every  $\ell \neq \ell^* \in L$ , either  $\ell$  and  $\ell^*$  are parallel, or there is exactly one  $p \in P$  such that  $pI\ell$  and  $pI\ell^*$ .
- (N) For every  $p \in P$  there is an  $\ell \in L$  such that  $p$  doesn't lie on  $\ell$  and for every  $\ell \in L$  there is an  $p \in P$  such that  $p$  doesn't lie on  $\ell$ .

Let  $\mathbf{P} := \langle \mathbb{R}^2, \mathcal{L}, \in \rangle$ . Then  $\mathbf{P}$  is a plane.

- (WE) (“the weak Euclidean postulate”) For every  $\ell \in L$  and every  $p \in P$  such that  $p$  doesn't lie on  $\ell$ , there is an  $\ell^* \in L$  such that  $pI\ell^*$  and  $\ell$  and  $\ell^*$  are parallel.
- (SE) (“the strong Euclidean postulate”) For every  $\ell \in L$  and every  $p \in P$  such that  $p$  doesn't lie on  $\ell$ , there is **exactly one**  $\ell^* \in L$  such that  $pI\ell^*$  and  $\ell$  and  $\ell^*$  are parallel.

$\mathbf{P}$  is a strongly Euclidean plane.

# Geometries (4).

**Question.** Do (A1), (A2), (N), and (WE) imply (SE)?

It is easy to see what a **positive solution** would be, but a **negative solution** would require reasoning over all possible proofs.

**Semantic version of the question.** Is every weakly Euclidean plane strongly Euclidean?

# Syntactic versus semantic.

	Does $\Phi$ imply $\psi$ ?	Does every $\Phi$ -structure satisfy $\psi$ ?
Positive	Give a proof $\exists$	Check all structures $\forall$
Negative	Check all proofs $\forall$	Give a counterexample $\exists$

# Euclid's Fifth Postulate.

“the scandal of elementary geometry” (D'Alembert 1767)

“In the theory of parallels we are even now not further than Euclid. This is a shameful part of mathematics...” (Gauss 1817)

Johann Carl Friedrich Gauss

(1777-1855)



1817

Nikolai Ivanovich Lobachevsky

(1792-1856)



1829

János Bolyai

(1802-1860)



1823

# A non-Euclidean geometry.

Take the usual geometry  $\mathbf{P} = \langle \mathbb{R}^2, \mathcal{L}, \in \rangle$  on the Euclidean plane.

Consider  $\mathbb{U} := \{x \in \mathbb{R}^2; \|x\| < 1\}$ . We define the restriction of  $\mathcal{L}$  to  $\mathbb{U}$  by  $\mathcal{L}^{\mathbb{U}} := \{\ell \cap \mathbb{U}; \ell \in \mathcal{L}\}$ .

$\mathbb{U} := \langle \mathbb{U}, \mathcal{L}^{\mathbb{U}}, \in \rangle$ .

**Theorem.**  $\mathbb{U}$  is a weakly Euclidean plane which is not strongly Euclidean.