

# A NOTE ON PERFECT MATCHINGS IN UNIFORM HYPERGRAPHS WITH LARGE MINIMUM COLLECTIVE DEGREE

VOJTĚCH RÖDL, ANDRZEJ RUCIŃSKI, MATHIAS SCHACHT, AND ENDRE SZEMERÉDI

ABSTRACT. For an integer  $k \geq 2$  and a  $k$ -uniform hypergraph  $H$ , let  $\delta_{k-1}(H)$  be the largest integer  $d$  such that every  $(k-1)$ -element set of vertices of  $H$  belongs to at least  $d$  edges of  $H$ . Further, let  $t(k, n)$  be the smallest integer  $t$  such that every  $k$ -uniform hypergraph on  $n$  vertices and with  $\delta_{k-1}(H) \geq t$  contains a perfect matching. The parameter  $t(k, n)$  has been completely determined for all  $k$  and large  $n$  divisible by  $k$  by Rödl, Ruciński, and Szemerédi in [*Perfect matchings in large uniform hypergraphs with large minimum collective degree*, submitted]. The values of  $t(k, n)$  are very close to  $n/2 - k$ . In fact, the function  $t(k, n) = n/2 - k + c_{n,k}$ , where  $c_{n,k} \in \{3/2, 2, 5/2, 3\}$  depends on the parity of  $k$  and  $n$ . The aim of this short note is to present a simple proof of an only slightly weaker bound:  $t(k, n) \leq n/2 + k/4$ . Our argument is based on an idea used in a recent paper of Aharoni, Georgakopoulos, and Sprüssel.

## 1. INTRODUCTION

A  $k$ -uniform hypergraph is a pair  $H = (V, E)$ , where  $V := V(H)$  is a finite set of vertices and  $E := E(H) \subseteq \binom{V}{k}$  is a family of  $k$ -element subsets of  $V$ . Whenever convenient we will identify  $H$  with  $E(H)$ . A *matching* in  $H$  is a set of disjoint edges of  $H$ .

Given a  $k$ -uniform hypergraph  $H$  and  $r$  vertices  $v_1, \dots, v_r \in V(H)$ ,  $1 \leq r \leq k-1$ , we denote by  $\deg_H(v_1, \dots, v_r)$  the number of edges of  $H$  which contain  $v_1, \dots, v_r$ . Let  $\delta_r(H) := \delta_r$  be the minimum of  $\deg_H(v_1, \dots, v_r)$  over all  $r$ -element sets of vertices of  $H$ .

**Definition 1.** For all integers  $k \geq 2$  and  $n \geq k$  divisible by  $k$ , denote by  $t(k, n)$  the smallest integer  $t$  such that every  $k$ -uniform hypergraph on  $n$  vertices and with  $\delta_{k-1} \geq t$  contains a perfect matching, that is, a matching of size  $n/k$ .

For graphs, an easy argument shows that  $t(2, n) = n/2$ . It follows from [3] that  $t(k, n) \leq n/2 + o(n)$ . In [2], Kühn and Osthus proved that  $t(k, n) \leq n/2 + 3k^2\sqrt{n \log n}$ . This was further improved in [5] to  $t(k, n) \leq n/2 + C \log n$ . Finally, the precise result was proved in [4], where it was shown that  $t(k, n) = n/2 - k + c_{n,k}$ , where  $c_{n,k} \in \{3/2, 2, 5/2, 3\}$  depends on the parity of  $k$  and  $n$ . The aim of this short note is to present a simple proof of an only slightly weaker bound.

**Theorem 2.** For all  $k \geq 3$  and  $n$  divisible by  $k$ ,  $t(k, n) \leq n/2 + k/4$ .

---

The first author was supported by NSF grant DMS 0300529.

The second author was supported by grant N201036 32/2546 and part of research was performed at Emory University, Atlanta.

The fourth author was supported by NSF grants DMS-0100784 and DMS-0603745.

Our argument is based on an idea used in a recent paper of Aharoni, Georgakopoulos, and Sprüssel [1]. Answering a question from [2], those authors proved in [1] a similar result for  $k$ -partite,  $k$ -uniform hypergraphs. Their result says that if  $V(H) = V_1 \cup \dots \cup V_k$ ,  $|V_1| = \dots = |V_k| = n$ , and for every  $(k-1)$ -tuple of vertices  $(v_1, \dots, v_{k-1}) \in V_1 \times \dots \times V_{k-1}$  we have  $\deg_H(v_1, \dots, v_{k-1}) > n/2$ , while for every  $(v_2, \dots, v_k) \in V_2 \times \dots \times V_k$  we have  $\deg_H(v_2, \dots, v_k) \geq n/2$ , then  $H$  has a perfect matching. While their simple and elegant approach does not seem to readily yield the precise function  $t(n, k)$ , it can be modified to prove Theorem 2.

## 2. PROOF OF THEOREM 2

Let  $H$  be a  $k$  uniform hypergraph on  $n$  vertices, where  $n$  is divisible by  $k$ , such that  $\delta_{k-1}(H) \geq n/2 + k/4$ . Further, let  $M$  be a largest matching in  $H$ . Suppose to the contrary that  $|M| \leq n/k - 1$ , that is,  $M$  is not perfect. By adding fake edges if necessary, without loss of generality we may assume that  $|M| = n/k - 1$ . (Alternatively, one could apply Proposition 2.1 from [4] – see Remark 2.1 there, which says that  $H$  contains a matching of size at least  $n/k - 1$ , if  $\delta_{k-1}(H) \geq n/k + O(\log n)$ .) Let  $x_1, \dots, x_k$  be the vertices of  $H$  not covered by  $M$ .

For every  $u \in V(M)$ , let  $e_u$  be the edge of  $M$  containing  $u$ . For every vertex  $v$  of  $H$ , let  $T_M(v)$  be the set of vertices  $u \in V(M)$  such that  $(e_u \setminus \{u\}) \cup \{v\}$  is an edge of  $H$ . Set  $t_M(v) = |T_M(v)|$ .

**Observation 1.** For each  $i = 1, \dots, k$ ,  $t_M(x_i) \leq n/2 - 5k/4$ .

*Proof.* If, say,  $t_M(x_k) > n/2 - 5k/4$ , then  $\deg_H(x_1, \dots, x_{k-1}) + t_M(x_k) > n - k = |V(M)|$ , so  $N(x_1, \dots, x_{k-1}) \cap T_M(x_k) \neq \emptyset$ . Let  $u \in N(x_1, \dots, x_{k-1}) \cap T_M(x_k)$ . Then, setting  $e' = \{u, x_1, \dots, x_{k-1}\}$  and  $e'' = (e_u \setminus \{u\}) \cup \{x_k\}$ , we see that  $M' = (M \setminus \{e_u\}) \cup \{e', e''\}$  is a perfect matching in  $H$  – a contradiction.  $\square$

**Observation 2.** There exists  $w \in V(M)$  with  $t_M(w) > n/2 - k/4$ .

*Proof.* Let  $B = (X \dot{\cup} Y, E_B)$  be an auxiliary bipartite graph where  $X = V(M)$ ,  $Y = V(H)$ , and  $uv \in E_B$  if and only if  $u \in X$ ,  $v \in Y$ , and  $u \in T_M(v)$ . In view of the assumption on  $\delta_{k-1}(H)$ , for each of the  $n - k$  vertices  $u \in X$  we have  $\deg_B(u) \geq n/2 + k/4$ . Let  $Y' = Y \setminus \{x_1, \dots, x_k\}$ . Then, in view of Observation 1, the number of edges in the induced subgraph  $B' = B[X \cup Y']$  is at least

$$(n - k) \left( \frac{n}{2} + \frac{k}{4} \right) - k \left( \frac{n}{2} - \frac{5k}{4} \right).$$

Hence, by averaging, there exists  $w \in Y' = V(M)$  such that

$$t_M(w) = \deg_{B'}(w) \geq \frac{e(B')}{n - k} \geq \left( \frac{n}{2} + \frac{k}{4} \right) - \frac{k(n/2 - 5k/4)}{n - k} > \frac{n}{2} - \frac{k}{4}.$$

$\square$

Fix  $w$  as in Observation 2.

**Observation 3.** There exists two vertices  $v_1$  and  $v_2$  and an edge  $e \in M \setminus \{e_w\}$  such that  $\{v_1, v_2\} \subseteq e$ ,  $v_1 \in N_H(e_w \setminus \{w\})$ , and  $v_2 \in N_H(x_1, \dots, x_{k-1})$ .

*Proof.* Together, the  $(k-1)$ -tuples  $S_1 = e_w \setminus \{w\}$  and  $S_2 = \{x_1, \dots, x_{k-1}\}$  have at most  $2(k+1) - 1 = 2k + 1$  neighbors in  $e_w \cup \{x_1, \dots, x_k\}$ . Thus, the total number

of pairs  $(v, i)$ , where  $v \in N_H(S_i)$ ,  $v \notin e_w \cup \{x_1, \dots, x_k\}$ , and  $i = 1, 2$ , is at least  $2(n/2 + k/4) - 2k - 1$ , and, by averaging, there exists  $e \in M \setminus \{e_w\}$  for which

$$|\{(v, i): v \in N_H(S_i) \cap e, i = 1, 2\}| \geq \frac{n + k/2 - 2k - 1}{n/k - 2} > k.$$

Consequently, there exist  $v_1, v_2 \in e$ ,  $v_1 \neq v_2$ , such that  $v_i \in N_H(S_i)$ ,  $i = 1, 2$ .  $\square$

By Observation 3, setting  $e' = (e_w \setminus \{w\}) \cup \{v_1\}$  and  $e'' = \{x_1, \dots, x_{k-1}, v_2\}$ , one can replace  $M$  with another matching  $M' = (M \setminus \{e_w, e\}) \cup \{e', e''\}$  of the same size, but such that  $w \notin V(M')$ . Note that  $T_M(w) \setminus T_{M'}(w) \subseteq e$ , and so,

$$t_{M'}(w) \geq t_M(w) - k > n/2 - 5k/4.$$

This is, however, a contradiction to Observation 1 (applied to  $M'$ ). This completes the proof of Theorem 2.

*Remark 3.* We believe that the bound on  $t(n, k)$  from Theorem 2 can be improved slightly, with a more cumbersome case analysis. However, for a clearer presentation we avoided those details.

**Acknowledgements.** We are grateful to the authors of [1] for providing us with an early version of their manuscript.

## REFERENCES

- [1] R. Aharoni, A. Georgakopoulos, and Ph. Sprüssel, *Perfect matchings in  $r$ -partite  $r$ -graphs*, submitted. [1](#), [2](#)
- [2] D. Kühn and D. Osthus, *Matchings in hypergraphs of large minimum degree*, J. Graph Theory **51** (2006), no. 4, 269–280. [1](#), [1](#)
- [3] V. Rödl, A. Ruciński, and E. Szemerédi, *An approximative Dirac-type theorem for  $k$ -uniform hypergraphs*, Combinatorica, to appear. [1](#)
- [4] ———, *Perfect matchings in large uniform hypergraphs with large minimum collective degree*, submitted. [1](#), [2](#)
- [5] ———, *Perfect matchings in uniform hypergraphs with large minimum degree*, European J. Combin. **27** (2006), no. 8, 1333–1349. [1](#)

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, GA 30322, USA

*E-mail address:* [rod1@mathcs.emory.edu](mailto:rod1@mathcs.emory.edu)

A. MICKIEWICZ UNIVERSITY, DEPARTMENT OF DISCRETE MATHEMATICS, UMULTOWSKA 87, 61-614 POZNAŃ, POLAND

*E-mail address:* [rucinski@amu.edu.pl](mailto:rucinski@amu.edu.pl)

INSTITUT FÜR INFORMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, UNTER DEN LINDEN 6, D-10099 BERLIN, GERMANY

*E-mail address:* [schacht@informatik.hu-berlin.de](mailto:schacht@informatik.hu-berlin.de)

DEPARTMENT OF COMPUTER SCIENCE, RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019, USA

*E-mail address:* [szemered@cs.rutgers.edu](mailto:szemered@cs.rutgers.edu)