

RAMSEY-TYPE NUMBERS INVOLVING GRAPHS AND HYPERGRAPHS WITH LARGE GIRTH

HIỆP HÀN, TROY RETTER, VOJTĚCH RÖDL, AND MATHIAS SCHACHT

ABSTRACT. Erdős asked if for every pair of positive integers r and k , there exists a graph H having $\text{girth}(H) = k$ and the property that every r -colouring of the edges of H yields a monochromatic cycle C_k . The existence of such graphs H was confirmed by the third author and Ruciński.

We consider the related numerical problem of estimating the order of the smallest graph H with this property for given integers r and k . We show that there exists a graph H on $R^{10k^2} k^{15k^3}$ vertices (where $R = R(C_k; r)$ is the r -colour Ramsey number for the cycle C_k) having $\text{girth}(H) = k$ and the Ramsey property that every r -colouring of the edges of H yields a monochromatic C_k . Two related numerical problems regarding arithmetic progressions in subsets of the integers and cliques in graphs are also considered.

§1. INTRODUCTION

For an integer $r \geq 2$ and graphs H and F , we write $H \rightarrow (F)_r$ if every r -colouring of the edges of H yields a monochromatic copy of F . If $H \rightarrow (F)_r$, we say that H is *Ramsey for F for r colours*. It follows from Ramsey's theorem that for every graph F and for every positive integer r , there exists a graph H such that $H \rightarrow (F)_r$. We consider three Ramsey-type problems that pertain to cycles in graphs and hypergraphs.

1.1. Cycles in Graphs. Our first result relates to a problem suggested by Erdős (see, e.g., [8]), which asks if for every pair of positive integers r and k , there exists a graph H having $\text{girth}(H)$ exactly k and the Ramsey property $H \rightarrow (C_k)_r$. The existence of such graphs was first established in [24]. We address the associated numerical problem.

Theorem 1.1. *Let $R = R(C_k; r)$ be the Ramsey number that denotes the least integer m such that $K_m \rightarrow (C_k)_r$. Then for all integers $k \geq 4$ and $r \geq 2$, there exists a graph H on $|V(H)| = k^{15k^3} R^{10k^2}$ vertices satisfying $\text{girth}(H) = k$ and $H \rightarrow (C_k)_r$.*

2010 *Mathematics Subject Classification.* 05C55 (primary), 05D10 (secondary).

Key words and phrases. Ramsey numbers, high girth.

H. Hàn was partly supported by FAPESP (2010/16526-3 and 2013/11353-1).

V. Rödl was supported by NSF grants DMS 1301698 and 1764385.

M. Schacht was supported through the Heisenberg-Programme of the DFG.

The exponential dependency of $|V(H)|$ on k in Theorem 1.1 is unavoidable as shown in the following Theorem 1.2. Further, note that the r -colour Ramsey number $R(C_k; r)$ is polynomial in r for fixed even k , while for fixed odd k it satisfies the exponential relation $c_1^r \leq R(C_k; r) \leq c_2^{r \log r}$ for some positive constants c_1 and c_2 (see, e.g., [10]). This leads to the following bounds, which show that the additional girth requirement on H still admits Ramsey graphs of order comparable to $R(C_k, r)$. We define

$$f(k, r) = \min \{|V(H)| : \text{girth}(H) = k \text{ and } H \longrightarrow (C_k)_r\}$$

and write $\log(x)$ for the binary logarithm and $\ln(x)$ for the natural logarithm.

Theorem 1.2. *There exist positive constants c_1 and c_2 such that for all $k \geq 2$ and $r \geq 2$*

$$\exp(c_1 k \log r) \leq f(2k, r) \leq \exp(c_2 k^2 (\log r + k \log k))$$

and

$$\exp(c_1 k r) \leq f(2k - 1, r) \leq \exp(c_2 k^2 (r \log r + k \log k)).$$

In particular, for fixed $k \geq 2$ we have $f(2k, r) = r^{\Theta(1)}$ and $e^{\Omega(r)} \leq f(2k - 1, r) \leq e^{O(r \log r)}$.

In Section 6 we present a simpler and more efficient argument for $f(6, r) = O(r^6)$, $f(8, r) = O(r^{12})$, and $f(12, r) = O(r^{30})$, which utilises known extremal constructions for graphs without short even cycles. Furthermore, we remark that the factor of $\log r$ in the upper bound on $f(2k - 1, r)$ can be removed for large k using the same proof together with the recent result from [17].

Proof. To show the upper bound for $f(2k, r)$ we note that it was announced in [7] and proven in [5] that, for every integer $k \geq 2$ every graph on n vertices with at least $100kn^{1+1/k}$ edges contains a copy of the cycle C_{2k} . For $n \geq 20kr^{k/(k-1)}$ we have $\binom{n}{2}/r \geq 100kn^{1+1/k}$ and consequently every edge colouring of K_n with r colours will have a monochromatic cycle C_{2k} . This shows that $R(C_{2k}; r) \leq 20kr^{k/(k-1)}$ and the bound $f(2k, r) \leq \exp(c_2 k^2 (\log r + k \log k))$ now follows from substituting into Theorem 1.1.

Similarly, we obtain the upper bound for $f(2k - 1, r)$ from Theorem 1.1 together with the bound $R(C_{2k-1}; r) \leq (r + 2)! \cdot (2k - 1)$ for $k \geq 2$ which was established in [4].

We now turn our attention towards the lower bounds. For any $k \geq 2$ and $r \geq 2$ consider any graph H with $\text{girth}(H) = 2k$ and the property $H \longrightarrow (C_{2k})_r$. Let $\tilde{H} \subseteq H$ be an edge minimal subgraph such that $\tilde{H} \longrightarrow (C_{2k})_r$. Clearly the minimum degree of \tilde{H} must be at least r and \tilde{H} must have girth $2k$. Since any graph with girth $2k$ and minimum degree r must have at least $2 \sum_{i=0}^{k-1} (r-1)^i \geq cr^{k-1}$ vertices the lower bound for $f(2k, r)$ follows.

To establish that $f(2k - 1, r) \geq \exp(c_1 k r)$ for any $k \geq 2$ and $r \geq 2$, as before we begin by considering any graph H with $\text{girth}(H) = 2k - 1$ and the property $H \longrightarrow (C_{2k-1})_r$.

Note that $\chi(H) > 2^r$, since otherwise the edges of H could be decomposed into r bipartite graphs, resulting in an r -colouring of $E(H)$ with no monochromatic odd cycle. Moreover, since $\chi(H) > 2^r$, there must be a subgraph $\tilde{H} \subset H$ with minimum degree at least 2^r . Since \tilde{H} has at least girth $2k - 1$ and minimum degree 2^r , the number of vertices in \tilde{H} must be at least $1 + 2^r \sum_{i=1}^{k-2} (2^r - 1)^i \geq 2^{crk}$ vertices for some $c > 0$. \square

1.2. Arithmetic Progressions. For a subset $S \subseteq \mathbb{N}$ and integers $k \geq 3$ and $r \geq 2$, we write $S \rightarrow (AP_k)_r$ to signify that every r -colouring of the integers in S yields a monochromatic arithmetic progression of length k . Van der Waerden’s theorem shows for all integers $k \geq 3$ and $r \geq 2$ that there is some integer N such that $[N] \rightarrow (AP_k)_r$, where we denote by $[N]$ the set of the first N positive integers $\{1, 2, \dots, N\}$. Several refinements of this well known theorem have been considered. One generalisation, suggested by Erdős [9], asks if for all $k \geq 3$ and $r \geq 2$, there exists an AP_{k+1} -free set $S \subseteq \mathbb{N}$ that has the property $S \rightarrow (AP_k)_r$, where a set is AP_{k+1} -free if it does not contain an arithmetic progression of length $k + 1$. This was answered independently by Spencer [29] and by Nešetřil and Rödl [20]. Moreover, Graham and Nešetřil [15] showed that there exist arbitrarily large AP_{k+1} -free sets S that have the property $S \rightarrow (AP_k)_r$ and are minimal in the sense that, for every $s \in S$, the subset $S' = S \setminus \{s\}$ fails to have the van der Waerden property, i.e., $S' \not\rightarrow (AP_k)_r$.

Furthermore, one may want to restrict the structure of the arithmetic progressions of length k in a set $S \subseteq \mathbb{N}$, but keep the van der Waerden property. That is, consider the *system of copies* of arithmetic progression of length k in S , which is the k -uniform hypergraph $(S, \binom{S}{AP_k})$ on the vertex set S with edge set $\binom{S}{AP_k}$ consisting of the k element subsets of S that form arithmetic progressions of length k . For a simpler notation, it will be convenient to identify this hypergraph just by its edge set. Moreover, we denote its chromatic number simply by $\chi(\binom{S}{AP_k})$ instead of $\chi((\binom{S}{AP_k}))$. Similarly, we suppress the outer pair of parentheses for other numerical hypergraph parameters as well.

Observe that $S \rightarrow (AP_k)_r$ if and only if the chromatic number satisfies $\chi(\binom{S}{AP_k}) > r$. Hence, van der Waerden’s theorem establishes that for fixed k , the chromatic number $\chi(\binom{[N]}{AP_k}) \rightarrow \infty$ as N tends to infinity. In view of the result of Erdős and Hajnal [11], which establishes the existence of hypergraphs having both large chromatic number and large girth, it is naturally to ask the following. Does for all $k, g \geq 3$, and $r \geq 2$ there exist a set $S \subseteq \mathbb{N}$ so that the hypergraph $(\binom{S}{AP_k})$ satisfies both the properties

- (P1) $\chi(\binom{S}{AP_k}) > r$ and
- (P2) $\text{girth}(\binom{S}{AP_k}) \geq g$?

As usual we say a k -uniform hypergraph has girth at least g if, for any integer h with $2 \leq h < g$, any subset of h edges span at least $(k-1)h+1$ vertices. In particular, $\text{girth}_{(AP_k)^S} \geq 3$ implies that no two arithmetic progressions can intersect in more than one point, which implies that S is AP_{k+1} -free. The existence of sets $S \subseteq \mathbb{N}$ satisfying properties (P1) and (P2) was established in [23] (see also [24]) and our next result gives an upper bound for the size of the smallest such set S .

Theorem 1.3. *Let $W = \text{vdW}(k, r)$ denote the least integer N such that $[N] \rightarrow (AP_k)_r$. Then for all integers $k, g \geq 3$, and $r \geq 2$, there exists a set $S \subseteq \mathbb{N}$ such that*

$$\chi_{(AP_k)^S} > r, \quad \text{girth}_{(AP_k)^S} \geq g, \quad \text{and} \quad |S| \leq k^{40k^2(k+g)} W^{12k(k+g)}.$$

To illustrate the result, consider the special case $k = 3$ for fixed $g \geq 3$. A result of Sanders [26] (see also [3]) implies that $\text{vdW}(3; r) \leq \exp(r^{1+o(1)})$, where the error term $o(1) \rightarrow 0$ as $r \rightarrow \infty$. Consequently, our result yields the existence of a set S of size at most $\exp(r^{1+o(1)})$ such that the properties $S \rightarrow (AP_3)_r$ and $\text{girth}_{(AP_3)^S} \geq g$ both hold. In other words, as in Theorem 1.1 the added girth condition does not essentially increase the best known upper bound in this case.

1.3. Cliques in Graphs. Another well known problem of Erdős and Hajnal [12] asked if, for every pair of positive integers k and r , there exists a K_{k+1} -free graph H such that $H \rightarrow (K_k)_r$. The case $r = 2$ was confirmed by Folkman [14], and the general case $r > 2$ was resolved by Nešetřil and Rödl [19]. Subsequently, Erdős [8] asked for a strengthened form of this result, namely the existence of a graph H with $H \rightarrow (K_k)_r$ in which no two copies of K_k share more than one edge, which was established in [21] (see also [22] for a generalisation from cliques K_k to arbitrary graphs).

As in the context of van der Waerden's theorem in Section 1.2, we may consider the structure of the cliques in H in more detail, i.e., we consider the *system of copies* of K_k in H , which is the $\binom{k}{2}$ -uniform hypergraph $(E(H), \binom{H}{K_k})$ having vertex set $E(H)$ and hyperedges corresponding to the edge sets of copies of K_k in H . As above we identify this hypergraph by its edge set $\binom{H}{K_k}$ and denote by $\chi_{\binom{H}{K_k}}$ and $\text{girth}_{\binom{H}{K_k}}$ its chromatic number and its girth. Again the statement $H \rightarrow (K_k)_r$ is equivalent to $\chi_{\binom{H}{K_k}} > r$ and the property that any two copies of K_k in H share at most one edge is equivalent to $\text{girth}_{\binom{H}{K_k}} \geq 3$. We give a new proof of the result from [21] that leads to a new upper bound on the size of the smallest such H .

Theorem 1.4. *Let $R = R(K_k; r)$ be the Ramsey number that denotes the least integer m such that $K_m \rightarrow (K_k)_r$. Then for all integers $k, g \geq 3$, and $r \geq 2$, there exists a graph H*

such that

$$\chi\left(\begin{matrix} H \\ K_k \end{matrix}\right) > r, \quad \text{girth}\left(\begin{matrix} H \\ K_k \end{matrix}\right) \geq g, \quad \text{and} \quad |V(H)| \leq k^{40gk^4} R^{40gk^2}.$$

By reversing the dependency between g and $|V(H)|$, we obtain the following corollary.

Corollary 1.5. *For all integers $k \geq 3$ and $r \geq 2$, there exist $c_{k,r} > 0$ and n_0 such that, for every integer $n \geq n_0$, there exists a graph H on n vertices satisfying both $H \rightarrow (K_k)_r$ and $\text{girth}\left(\begin{matrix} H \\ K_k \end{matrix}\right) \geq c_{k,r} \log n$.*

It can be shown that any graph H on n vertices satisfying $H \rightarrow (K_k)_r$ must also satisfy $\text{girth}\left(\begin{matrix} H \\ K_k \end{matrix}\right) = O(\log n)$, due to the degree condition required by $\chi\left(\begin{matrix} H \\ K_k \end{matrix}\right) > r$ and, in that sense, our result gives an optimal order of magnitude for the girth.

1.4. Overview. In the proof of Theorems 1.1, 1.3, and 1.4, we use the *Container Method* from [1]. Although our three proofs are similar in nature, we demonstrate two slightly different approaches. In the proof of Theorem 1.1 we will use the container method and the FKG-inequality. We show that for suitable choice of parameters p and n , the probability that the random graph $G(n, p)$ fails to have the Ramsey-property for C_k and r is strictly smaller than the probability that it has girth at least k . This implies the existence of graphs which has both, the Ramsey property and large girth.

In contrast, for Theorems 1.3 and 1.4, we will first find objects which are robust with respect to the Ramsey property and which contain few short cycles. The desired objects are then obtained by a deletion argument.

In Section 2 we state the Container Lemma [1], which plays a central rôle in our proofs. The details of the proofs of Theorems 1.1, 1.3, and 1.4 are given in Sections 3, 4, and 5 respectively. Section 6 contains some additional remarks related to Theorem 1.1.

Acknowledgements. We are thankful for the thoughtful comments we received through the reviewing process.

§2. HYPERGRAPH CONTAINERS

The proofs of the theorems presented in Section 1 use many ideas from [18, 25] and rely on random constructions combined with the *Container Method* of Balogh, Morris, and Samotij [1] and of Saxton and Thomason [27]. For the numerical aspects the container result from [27] seemed to be better suited and we state it below (see Theorem 2.1).

Roughly speaking, this lemma states that, if a given hypergraph \mathcal{H} satisfies some numerical ‘degree conditions’, then there there exists a relatively ‘small’ family of sets of

so-called ‘containers’ (sets \mathcal{C} in Theorem 2.1 below) that are ‘almost’ independent sets of vertices that cover all independent sets of \mathcal{H} .

We now introduce the notation necessary for the formulation of this theorem. For a h -uniform hypergraph \mathcal{H} , let $e(\mathcal{H})$ denote the number of (hyper)edges in \mathcal{H} . For a set $J \subseteq V(\mathcal{H})$ we define the *degree of J* by

$$d(J) = |\{e \in E(\mathcal{H}) : e \supseteq J\}|$$

and for $j = 1, \dots, h$ we define the *maximum j -degree* of a vertex $v \in V(\mathcal{H})$ by

$$d_j(v) = \max \left\{ d(J) : J \in \binom{V(\mathcal{H})}{j} \text{ and } v \in J \right\}.$$

The average of $d_j(v)$ is denoted by

$$d_j = \frac{1}{|V(\mathcal{H})|} \sum_{v \in V} d_j(v).$$

Note that $d_1(v)$ is just the degree of v in \mathcal{H} and, consequently, d_1 denotes the average vertex degree of \mathcal{H} . With this notation at hand we state the *Container Lemma* from [27, Corollary 3.6].

Theorem 2.1 (Container Lemma). *Let $\mathcal{H} = (V, E)$ be a h -uniform hypergraph and suppose $\tau, \varepsilon \in (0, 1/2)$ satisfy*

$$\frac{6 \cdot h! \cdot 2^{\binom{h}{2}}}{d_1} \sum_{j=2}^h \frac{d_j}{2^{\binom{j-1}{2}} \tau^{j-1}} \leq \varepsilon. \quad (1)$$

Then for integers

$$K = 800h(h!)^3 \quad \text{and} \quad s = \lfloor K \log(1/\varepsilon) \rfloor$$

the following holds.

For every independent set $I \subseteq V$ in \mathcal{H} , there exists an s -tuple $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_s)$ of subsets of V and a subset $\mathcal{C} = \mathcal{C}(\mathcal{S}) \subseteq V$ only depending on \mathcal{S} such that

- (i) $\bigcup_{i \in [s]} \mathcal{S}_i \subseteq I \subseteq \mathcal{C}$,
- (ii) $e(\mathcal{H}[\mathcal{C}]) \leq \varepsilon \cdot e(\mathcal{H})$, and
- (iii) for every $i \in [s]$ we have $|\mathcal{S}_i| \leq \tau K |V|$.

The Container Lemma stated here is an abridged version of [27, Corollary 3.6], which suffices for our purpose. For the explicit constant 800 appearing in the constant K see the discussion following Corollary 3.6 in [27].

§3. PROOF OF THEOREM 1.1

In this section we prove Theorem 1.1. For that we have to show that there exists a graph H on at most $k^{15k^3}R^{10k^2}$ vertices (where $R = R(C_k; r)$ is the r -colour Ramsey number for C_k) with $\text{girth}(H) = k$ and the Ramsey property $H \longrightarrow (C_k)_r$.

Proof. For integers $k \geq 4$ and $r \geq 2$ let $R = R(C_k; r)$ be the r -colour Ramsey number for C_k . We first define all constants involved in the proof. For the application of the Container Lemma we set

$$\varepsilon = \frac{1}{rR^k} \quad \text{and} \quad D_\tau = \frac{2^{2k}}{\varepsilon^{1/(k-1)}} \quad (2)$$

and we fix integers

$$K = 800k(k!)^3 < 30k^{3k} \quad \text{and} \quad s = \lfloor K \log(1/\varepsilon) \rfloor < 30k^{3k} \log(rR^k). \quad (3)$$

We set

$$D_p = 10R^2r^2s^2KD_\tau \log(10R^2r) < k^{15k}R^{10} \quad (4)$$

and define

$$n = D_p^{k^2} < k^{15k^3}R^{10k^2}. \quad (5)$$

Finally, we define the following parameters appearing in the proof

$$\tau = D_\tau n^{-\frac{k-2}{k-1}} \quad \text{and} \quad p = D_p n^{-\frac{k-2}{k-1}}. \quad (6)$$

In the proof we consider the binomial random graph $G(n, p)$ and the theorem is a consequence of the following two claims, which we verify below.

Claim 3.1. $\mathbb{P}(\text{girth}(G(n, p)) \geq k) \geq \exp(-kn^{k-1}p^{k-1})$.

Claim 3.2. $\mathbb{P}(G(n, p) \longrightarrow (C_k)_r) \geq 1 - \exp(-\frac{p}{2R^2} \binom{n}{2})$.

In order to compare the probabilities in the above two claims, we observe

$$\frac{kn^{k-1}p^{k-1}}{p \binom{n}{2} / 2R^2} = \frac{4R^2kD_p^{k-1}}{p(n-1)} < \frac{8R^2kD_p^{k-1}}{pn} = \frac{8R^2kD_p^{k-1}}{D_p n^{1/(k-1)}} = \frac{8R^2k}{D_p^{3+1/(k-1)}} < 1.$$

With this in mind, it now follows from Claims 3.1 and 3.2 that with positive probability the random graph $G(n, p)$ has girth at least k and the Ramsey-property for C_k and r colours. Consequently, Theorem 1.1 follows from our choice of n in (5). \square

Proof of Claim 3.1. To bound of the probability that $G(n, p)$ has girth at least k , we first note that the number of cycles of length j in K_n is $\frac{(j-1)!}{2} \binom{n}{j}$, and that each cycle C_j will occur in $G(n, p)$ with probability p^j . To compute the probability that none of these (dependent) events occur, we use the FKG-inequality (see, e.g., [16, Section 2.2]). For this

purpose, let X_{k-1} be the random variable which counts the number of cycles of length less than k . Clearly, X_{k-1} is the sum of monotone increasing indicator random variables and, hence, the FKG-inequality asserts

$$\mathbb{P}(\text{girth}(G(n, p)) \geq k) = \mathbb{P}(X_{k-1} = 0) \stackrel{\text{FKG}}{\geq} \prod_{j=3}^{k-1} (1 - p^j)^{\frac{(j-1)!}{2} \binom{n}{j}} \geq \exp\left(-\frac{\mathbb{E}[X_{k-1}]}{1 - p^3}\right), \quad (7)$$

where we used the estimate $1 - x \geq \exp(-x/(1 - x))$ for the last inequality. Since $pn \geq 1$ we have

$$\mathbb{E}[X_{k-1}] = \sum_{j=3}^{k-1} \frac{(j-1)!}{2} \binom{n}{j} p^j \leq \sum_{j=3}^{k-1} \frac{(pn)^j}{2j} \leq \frac{k}{6} (pn)^{k-1} \stackrel{(6)}{=} \frac{k}{6} D_p^{k-1} n$$

and the claim follows from $1 - p^3 > 1/6$ and (7). \square

Proof of Claim 3.2. The proof relies on an application of the Container Lemma (Theorem 2.1) (for similar proofs see, e.g., [18, 25]). We will apply the Container Lemma to the k -uniform hypergraph $\mathcal{H} = \binom{K_n}{C_k}$ which is the system of all cycles C_k of length k in K_n , i.e., $V(\mathcal{H}) = E(K_n)$ and k edges of K_n correspond to a hyperedge in \mathcal{H} , if they form a cycle of length k . For the application of Theorem 2.1 we first verify (1). In that direction we note

$$d_1 = \frac{k \cdot |E(\mathcal{H})|}{|V(\mathcal{H})|} = \frac{k! \binom{n}{k}}{2 \binom{n}{2}} \geq \frac{k!}{k^k} n^{k-2}, \quad d_j \leq n^{k-j-1}$$

for $j = 2, \dots, k-1$, and $d_k = 1$. Therefore, for $j = 2, \dots, k-1$ we have

$$\frac{d_j}{d_1 \cdot \tau^{j-1}} \leq \frac{k^k}{k! \cdot (\tau n)^{j-1}} \leq \frac{k^k}{k! \cdot \tau n} = \frac{k^k}{k! \cdot D_\tau n^{1/(k-1)}} \leq \frac{k^k}{k! \cdot n^{1/(k-1)}}$$

and, moreover,

$$\frac{d_k}{d_1 \cdot \tau^{k-1}} \leq \frac{k^k}{k! n^{k-2} \cdot \tau^{k-1}} = \frac{k^k}{k! \cdot D_\tau^{k-1}}.$$

Combining both estimates yields

$$\frac{6 \cdot k! \cdot 2 \binom{k}{2}}{d_1} \sum_{j=2}^k \frac{d_j}{2 \binom{j-1}{2} \tau^{j-1}} \leq \frac{6 \cdot k^{k+1} \cdot 2 \binom{k}{2}}{\min\{n^{1/(k-1)}, D_\tau^{k-1}\}} \stackrel{(2),(5)}{\leq} \varepsilon.$$

Having verified (1) of Theorem 2.1, we infer properties (i)–(iii) for every independent set $I \subseteq V(\mathcal{H})$. We consider the family \mathcal{B} of all graphs $B \subseteq K_n$ that fail to have the Ramsey property, i.e., $B \not\rightarrow (C_k)_r$. Below we establish Claim 3.2 by showing

$$\mathbb{P}(G(n, p) \in \mathcal{B}) \leq \exp\left(-\frac{p}{2R^2} \binom{n}{2}\right).$$

By the definition of \mathcal{B} , for every $B \in \mathcal{B}$ there exists a partition $E(B) = I_1^B \cup \dots \cup I_r^B$ with the property that none of the sets I_i^B contains a cycle C_k . In particular, each I_i^B is an independent set in \mathcal{H} and, therefore, properties (i)–(iii) of the Container Lemma assert

that for every $i \in [r]$ there exists an s -tuple $\mathcal{S}_i^B = (\mathcal{S}_{i,1}^B, \dots, \mathcal{S}_{i,s}^B)$ of subsets of I_i^B and a container set $\mathcal{C}(\mathcal{S}_i^B) \supseteq I_i^B$ such that

$$|\mathcal{S}_{i,\sigma}^B| \leq \tau K \binom{n}{2}$$

for every $\sigma \in [s]$ and

$$|e(\mathcal{C}(\mathcal{S}_i^B))| \leq \varepsilon \cdot e(\mathcal{H}) = \varepsilon \cdot \frac{(k-1)!}{2} \binom{n}{k}. \quad (8)$$

We also set $\mathcal{S}^B = (\mathcal{S}_1^B, \dots, \mathcal{S}_r^B)$ and $\mathcal{C}^B = (\mathcal{C}(\mathcal{S}_1^B), \dots, \mathcal{C}(\mathcal{S}_r^B))$.

More generally, for any such possible r -tuple $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ of s -tuples of sets of size at most $\tau K \binom{n}{2}$, we consider the corresponding container vector $\mathcal{C}(\mathcal{S}) = (\mathcal{C}(\mathcal{S}_1), \dots, \mathcal{C}(\mathcal{S}_r))$ given by the Container Lemma. We denote by $D(\mathcal{S})$ its complement in $E(K_n)$ given by

$$D(\mathcal{S}) = E(K_n) \setminus (\mathcal{C}(\mathcal{S}_1) \cup \dots \cup \mathcal{C}(\mathcal{S}_r)).$$

We observe that for any $B \in \mathcal{B}$ the following two properties hold:

- (a) $\bigcup_{i \in [r]} \bigcup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}^B \subseteq E(B)$ and
- (b) $E(B) \cap D(\mathcal{S}^B) = \emptyset$, since

$$E(B) = I_1^B \cup \dots \cup I_r^B \subseteq \mathcal{C}(\mathcal{S}_1^B) \cup \dots \cup \mathcal{C}(\mathcal{S}_r^B) = E(K_n) \setminus D(\mathcal{S}^B).$$

From (a) and (b) we infer that

$$\begin{aligned} \mathbb{P}(G(n, p) \in \mathcal{B}) &\leq \sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} \mathbb{P}\left(\mathcal{S} \subseteq G(n, p) \wedge E(G(n, p)) \cap D(\mathcal{S}) = \emptyset\right) \\ &\leq \sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\bigcup_{i \in [r]} \bigcup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} \cdot \mathbb{P}\left(E(G(n, p)) \cap D(\mathcal{S}) = \emptyset\right) \\ &\leq \max_{\mathcal{S}} \mathbb{P}\left(E(G(n, p)) \cap D(\mathcal{S}) = \emptyset\right) \sum_{\mathcal{S}} p^{|\bigcup_{i \in [r]} \bigcup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|}, \end{aligned} \quad (9)$$

where the sum and the maximum are taken over all r -tuples $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ of s -tuples $\mathcal{S}_i = (\mathcal{S}_{i,1}, \dots, \mathcal{S}_{i,s})$ of sets $\mathcal{S}_{i,\sigma}$ of size at most $\tau K \binom{n}{2}$ for $i \in [r]$ and $\sigma \in [s]$. We will use property (ii) of the Container Lemma to bound the maximum probability, while our choice of constants allow us to derive a sufficient bound for the sum.

For the maximum probability below we first observe that for every $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ we have

$$|D(\mathcal{S})| > \frac{1}{R^2} \binom{n}{2}. \quad (10)$$

For the proof we use the fact that for any $(r+1)$ -colouring of $E(K_n)$ either there are more than

$$\frac{1}{2^{\binom{R}{k}}} \binom{n}{k}$$

monochromatic copies of C_k in the first r colours or there are more than $\frac{1}{R^2} \binom{n}{2}$ edges having the last colour (see, e.g., [25, Proposition 8] for the same assertion for cliques K_k instead of cycles and Fact 4.3 for the analogous statement for arithmetic progressions and its proof).

In view of this fact, we consider $\mathcal{C}(\mathcal{S}_1) \cup \dots \cup \mathcal{C}(\mathcal{S}_r) \cup D(\mathcal{S})$ as an $(r+1)$ -colouring of $E(K_n)$. Owing to property (ii) of the Container Lemma (see (8)), every $\mathcal{C}(\mathcal{S}_i)$ contains at most $\varepsilon \frac{(k-1)!}{2} \binom{n}{k}$ copies of C_k and, hence, there are at most

$$r \cdot \varepsilon \frac{(k-1)!}{2} \binom{n}{k} \stackrel{(2)}{\leq} \frac{1}{2 \binom{R}{k}} \binom{n}{k}$$

monochromatic copies in the first r colours. (We remark that we chose ε to satisfy the above inequality; in fact we took ε slightly smaller to make the calculations simpler.) Therefore, the mentioned fact above yields (10) and, consequently, we arrive at

$$\mathbb{P}\left(E(G(n, p)) \cap D(\mathcal{S}) = \emptyset\right) = (1-p)^{|D(\mathcal{S})|} \stackrel{(10)}{\leq} \exp\left(-\frac{p}{R^2} \binom{n}{2}\right) \quad (11)$$

for every \mathcal{S} considered here. In particular, (11) bounds the maximum probability considered in the R-H-S of (9) and below we turn to the sum in (9).

Owing to $|\mathcal{S}_{i,\sigma}| \leq \tau K \binom{n}{2}$ for every $i \in [r]$ and $\sigma \in [s]$ we have

$$\sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} \leq \sum_{m=0}^{r \cdot s \cdot \tau K \binom{n}{2}} \binom{\binom{n}{2}}{m} 2^{rsm} p^m \leq \sum_{m=1}^{r \cdot s \cdot \tau K \binom{n}{2}} \left(\frac{e \binom{n}{2}}{m} 2^{rs} p\right)^m.$$

Since the function $m \mapsto (e \binom{n}{2} 2^{rs} p / m)^m$ is unimodal and attains its maximum value for $m_0 = \binom{n}{2} 2^{rs} p \geq r s \tau K \binom{n}{2}$, we can bound the summands in R-H-S above by the last one and obtain

$$\begin{aligned} \sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} &\leq \left(r s \tau K \binom{n}{2} + 1\right) \cdot \left(\frac{e \binom{n}{2}}{r s \tau K \binom{n}{2}} 2^{rs} p\right)^{r s \tau K \binom{n}{2}} \\ &\leq n^2 \cdot \left(\frac{2^{rs} e D_p}{r s K D_\tau}\right)^{r s \tau K \binom{n}{2}} \\ &\stackrel{(4)}{=} n^2 \cdot (2^{rs} e \cdot 10R^2 r s \log(10R^2 r))^{r s \tau K \binom{n}{2}} \\ &\leq n^2 \cdot \exp\left(r s \tau K \binom{n}{2} (rs + 1 + \ln(10R^2 r s \log(10R^2 r)))\right) \\ &\stackrel{(4)}{\leq} n^2 \cdot \exp\left(\frac{p}{3R^2} \binom{n}{2}\right) \\ &\leq \exp\left(\frac{p}{2R^2} \binom{n}{2}\right). \end{aligned} \quad (12)$$

Finally, combining (11) and (12) with (9) leads to

$$\mathbb{P}(G(n, p) \in \mathcal{B}) \leq \exp\left(-\frac{p}{R^2} \binom{n}{2}\right) \cdot \exp\left(\frac{p}{2R^2} \binom{n}{2}\right) = \exp\left(-\frac{p}{2R^2} \binom{n}{2}\right),$$

which concludes the proof of the claim. \square

§4. PROOF OF THEOREM 1.3

We prove Theorem 1.3 by establishing that, for given integers $k \geq 3$, $r \geq 2$, and $g \geq 2$, there exists a set $S \subseteq \mathbb{N}$ of size at most $k^{40k^2(k+g)}W^{12k(k+g)}$ (where $W = \text{vdW}(k; r)$ is the van der Waerden number guaranteeing monochromatic arithmetic progressions of length k for any r -colouring of $[W]$) such that the hypergraph $(\binom{S}{AP_k})$ has chromatic number greater than r , and girth at least g .

Proof. Let $k \geq 3$, $r \geq 2$, and $g \geq 2$ be given and let $W = \text{vdW}(k; r)$ be the van der Waerden number. We first define all constants involved in the proof. For the application of the Container Lemma we set

$$\varepsilon = \frac{1}{rW^3} \quad \text{and} \quad D_\tau = \left(\frac{6 \cdot k! \cdot 2^{\binom{k}{2}} \cdot k^3}{\varepsilon}\right)^{\frac{1}{k-1}} \quad (13)$$

and we fix integers

$$K = 800k(k!)^3 < 30k^{3k} \quad \text{and} \quad s = \lfloor K \log(1/\varepsilon) \rfloor < 30k^{3k} \log(rW^3). \quad (14)$$

We set

$$D_p = 128Wr^2s^2KD_\tau \log(128Wr) < 2^{40}k^{10k}r^3W^3 \quad (15)$$

and define

$$n = k^{4g}D_p^{2k(k+g)} < k^{40k^2(k+g)}W^{12k(k+g)}. \quad (16)$$

Finally, we define the following parameters appearing in the proof

$$\tau = D_\tau n^{-\frac{1}{k-1}}, \quad p = D_p n^{-\frac{1}{k-1}}, \quad \text{and} \quad t = \frac{pn}{8W}. \quad (17)$$

Let $[n]_p$ denote the random set obtained by choosing each element of $[n] = \{1, 2, \dots, n\}$ independently with probability p . The theorem is an immediate consequence of the following two claims.

Claim 4.1. *With probability larger than $1/2$ for the random subset $[n]_p$ there exists a set $T \subseteq [n]_p$ of size at most t such that $\text{girth}(\binom{[n]_p \setminus T}{AP_k}) \geq g$.*

Claim 4.2. *With probability larger than $1/2$ the random subset $[n]_p$ satisfies $\chi(\binom{[n]_p \setminus T}{AP_k}) > r$ for every subset $T \subseteq [n]_p$ of size at most t .*

Together, these claims establish that, with positive probability, the random set $[n]_p$ will have the property that there exists a set $T \subseteq [n]_p$ of size t so that the hypergraph $\binom{[n]_p \setminus T}{AP_k}$ has girth at least g and chromatic number bigger than r . Thus, these claims together with our choice of n in (16) establish the existence of a set $S \subseteq \mathbb{N}$ as claimed in Theorem 1.3. We remark that, such a set will likely have only $O(pn)$ elements (not n elements). However, this improvement is negligible. \square

Proof of Claim 4.1. The proof follows by a standard first moment argument. Recall that a 2-cycle in a hypergraph consists of two hyperedges sharing at least two vertices and for $j > 2$ a j -cycle consists of a cyclically ordered sequence of hyperedges e_1, e_2, \dots, e_j where the intersection of two consecutive edges is exactly 1, the intersection of any two nonconsecutive edges is empty, and the intersection points for each pair of consecutive edges is unique (which for $j \geq 4$ is already implied by the other two conditions). Let the random variable X_j denote the number of j -cycles appearing in the random hypergraph $\binom{[n]_p}{AP_k}$.

We first estimate $\mathbb{E}[X_2]$. Since the hyperedges of $\binom{[n]}{AP_k}$ are arithmetic progressions of length k , every pair of vertices is contained in at most $\binom{k}{2}$ such hyperedges. Consequently, we have

$$\mathbb{E}[X_2] \leq \binom{n}{2} \binom{k}{2}^2 p^{k+1} \leq k^4 p^{k+1} n^2 = pn \cdot \frac{k^4 D_p^k}{n^{1/(k-1)}} < \frac{t}{4}.$$

Next we bound $\mathbb{E}[X_j]$ for $3 \leq j < g$. For that we note that for any j -cycle we may first select and order the j vertices of degree 2 and then fixing the remaining vertices of each edge. However, since every edge of the cycle contains two (already fixed) vertices of degree two, again there are at most $\binom{k}{2}$ possible completion for such an edge and, hence, we have

$$\sum_{j=3}^{g-1} \mathbb{E}[X_j] \leq \sum_{j=3}^{g-1} n^j k^{2j} p^{(k-1)j} = \sum_{j=3}^{g-1} k^{2j} D_p^{(k-1)j} < k^{2g} D_p^{kg} \leq \frac{t}{4}.$$

By Markov's inequality this implies that with probability less than $1/2$ the randomly generated hypergraph $\binom{[n]_p}{AP_k}$ has at most t cycles of length less than g , which establishes Claim 4.1. \square

Proof of Claim 4.2. We consider the k -uniform hypergraph $\mathcal{H} = \binom{[n]}{AP_k}$ and check that it satisfies the assumptions of the Container Lemma (Theorem 2.1) for the parameters ε and τ chosen in (13) and (17). Note that $\varepsilon < 1/2$ by definition and $\tau < 1/2$ follows from the choice of n in (16). For the remaining assumption (1) we recall the definition of the average degrees d_j for $j = 1, \dots, k$ of \mathcal{H} and again, using the fact that every pair of vertices is contained in at most $\binom{k}{2}$ AP_k 's, we note that for $j = 2, \dots, k$ we have

$$d_j \leq d_2 \leq \binom{k}{2} < \frac{k^2}{2}.$$

Moreover, we have

$$d_1 = \frac{k}{n} \cdot \left| \binom{[n]}{AP_k} \right| = \frac{k}{n} \cdot \sum_{i=1}^{n-k+1} \binom{n-i}{k-1} = \frac{k}{n} \cdot \sum_{i=1}^{n-1} \binom{n-i}{k-1} \geq \frac{k}{n} \cdot \sum_{i=1}^{n-1} \left(\frac{n-i}{k-1} - 1 \right) \geq \frac{n}{2}. \quad (18)$$

Consequently,

$$\frac{6 \cdot k! \cdot 2^{\binom{k}{2}}}{d_1} \sum_{j=2}^k \frac{d_j}{2^{\binom{j-1}{2}} \tau^{j-1}} \leq \sum_{j=2}^k \frac{6 \cdot k! \cdot 2^{\binom{k}{2}} \cdot k^2}{n \cdot 2^{\binom{j-1}{2}} \cdot D_\tau^{j-1} n^{-\frac{j-1}{k-1}}} < \frac{6 \cdot k! \cdot 2^{\binom{k}{2}} \cdot k^3}{\min\{D_\tau^{k-1}, n^{\frac{1}{k-1}}\}} \stackrel{(13),(16)}{\leq} \varepsilon.$$

This shows that condition (1) of Theorem 2.1 holds. Consequently, for every independent set $I \subseteq V(\mathcal{H})$ we can apply conclusions (i)–(iii) of the Container Lemma with the constants defined in (14).

We consider the family \mathcal{B} of all sets $B \subseteq [n]$ with the property that there exists a set $T \subseteq E(B)$ of size at most t such that $(B \setminus T) \dashrightarrow (AP_k)_r$, i.e., $\chi_{AP_k}^{(B \setminus T)} \leq r$. Claim 4.2 is equivalent to

$$\mathbb{P}([n]_p \in \mathcal{B}) < \frac{1}{2}.$$

By definition of \mathcal{B} , for every $B \in \mathcal{B}$ there exists a set $T^B \subseteq B$ of size $|T^B| \leq t$ and a partition $B = I_1^B \cup \dots \cup I_r^B \cup T^B$ with the property that none of the sets I_i^B contains an AP_k . In particular, each I_i^B is an independent set in \mathcal{H} and, therefore, properties (i)–(iii) of the Container Lemma assert that for every $i \in [r]$ there exists an s -tuple $\mathcal{S}_i^B = (\mathcal{S}_{i,1}^B, \dots, \mathcal{S}_{i,s}^B)$ of subsets of I_i^B and a container set $\mathcal{C}(\mathcal{S}_i^B) \supseteq I_i^B$ such that

$$|\mathcal{S}_{i,\sigma}^B| \leq \tau K n \quad (19)$$

for every $\sigma \in [s]$ and

$$|e_{\mathcal{H}}(\mathcal{C}(\mathcal{S}_i^B))| \leq \varepsilon \cdot e(\mathcal{H}) = \varepsilon \cdot \left| \binom{[n]}{AP_k} \right|. \quad (20)$$

We set $\mathcal{S}^B = (\mathcal{S}_1^B, \dots, \mathcal{S}_r^B)$ and $\mathcal{C}^B = (\mathcal{C}(\mathcal{S}_1^B), \dots, \mathcal{C}(\mathcal{S}_r^B))$.

Moreover, for any possible r -tuple $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ of s -tuples of sets of size at most $\tau K n$, we consider the corresponding container vector $\mathcal{C}(\mathcal{S}) = (\mathcal{C}(\mathcal{S}_1), \dots, \mathcal{C}(\mathcal{S}_r))$ given by the Container Lemma. We denote by $D(\mathcal{S})$ its complement in $[n]$ given by

$$D(\mathcal{S}) = [n] \setminus (\mathcal{C}(\mathcal{S}_1) \cup \dots \cup \mathcal{C}(\mathcal{S}_r)).$$

Observe that for any $B \in \mathcal{B}$ the following two properties hold:

- (a) $\bigcup_{i \in [r]} \bigcup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}^B \subseteq (E(B) \setminus D(\mathcal{S}^B))$ and
- (b) $|B \cap D(\mathcal{S}^B)| \leq |T^B| \leq t$, since

$$B \setminus T^B = I_1^B \cup \dots \cup I_r^B \subseteq \mathcal{C}(\mathcal{S}_1^B) \cup \dots \cup \mathcal{C}(\mathcal{S}_r^B) = [n] \setminus D(\mathcal{S}^B).$$

From (a) and (b) we infer that

$$\begin{aligned} \mathbb{P}([n]_p \in \mathcal{B}) &\leq \sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} \cdot \mathbb{P}(|[n]_p \cap D(\mathcal{S})| \leq t) \\ &\leq \max_{\mathcal{S}} \mathbb{P}(|[n]_p \cap D(\mathcal{S})| \leq t) \cdot \sum_{\mathcal{S}} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|}, \end{aligned} \quad (21)$$

where the sum and the maximum are taken over all r -tuples $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ of s -tuples $\mathcal{S}_i = (\mathcal{S}_{i,1}, \dots, \mathcal{S}_{i,s})$ of sets $\mathcal{S}_{i,\sigma}$ of size at most τKn for $i \in [r]$ and $\sigma \in [s]$. We will use property (ii) of the Container Lemma to bound the maximum probability in (21), while our choice of constants allow us to derive a sufficient bound for the sum in (21). We shall use the following fact (the proof of which we defer to the end of this section).

Fact 4.3. *For every $(r+1)$ -colouring of $[n]$ either there are more than $\lfloor \binom{[n]}{AP_k} \rfloor / W^3$ monochromatic AP_k 's in the first r colours or more than $\frac{n}{4W}$ elements are in the last colour.*

In view of this fact, we consider $\mathcal{C}(\mathcal{S}_1) \cup \dots \cup \mathcal{C}(\mathcal{S}_r) \cup D(\mathcal{S})$ as an $(r+1)$ -colouring of $[n]$. Owing to property (ii) of the Container Lemma (see (20)), every $\mathcal{C}(\mathcal{S}_i)$ contains at most $\varepsilon \lfloor \binom{[n]}{AP_k} \rfloor$ monochromatic AP_k 's and, hence, there are at most

$$r \cdot \varepsilon \left| \binom{[n]}{AP_k} \right| \stackrel{(13)}{=} \frac{1}{W^3} \left| \binom{[n]}{AP_k} \right|$$

monochromatic AP_k 's in the first r colours. Therefore, Fact 4.3 yields that for every $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ we have

$$|D(\mathcal{S})| > \frac{n}{4W}. \quad (22)$$

In particular, the choice of t combined with (22) yields $t < p|D(\mathcal{S})|/2$ and, consequently, Chernoff's inequality (see, e.g., [16, Theorem 2.1] asserts

$$\mathbb{P}(|[n]_p \cap D(\mathcal{S})| \leq t) \leq \exp\left(-\frac{pn}{32W}\right) \quad (23)$$

for every \mathcal{S} considered here. In particular, (23) bounds the maximum probability considered in the R-H-S of (21) and below we turn to the sum in (21).

Owing to $|\mathcal{S}_{i,\sigma}| \leq \tau Kn$ for every $i \in [r]$ and $\sigma \in [s]$ we have

$$\sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} \leq \sum_{m=0}^{r \cdot s \cdot \tau Kn} \binom{n}{m} 2^{rsm} p^m \leq \sum_{m=0}^{r \cdot s \cdot \tau Kn} \left(\frac{en}{m} 2^{rs} p\right)^m. \quad (24)$$

Since the function $m \mapsto (en2^{rs}p/m)^m$ is unimodal and attains its maximum value for $m_0 = 2^{rs}pn \geq rs\tau Kn$, from (24) we obtain

$$\begin{aligned}
\sum_{\mathcal{S}=(\mathcal{S}_1,\dots,\mathcal{S}_r)} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} &\leq (rs\tau Kn + 1) \cdot \left(\frac{en}{rs\tau Kn} 2^{rs}p \right)^{rs\tau Kn} \\
&\leq n \cdot \left(\frac{2^{rs}eD_p}{rsKD_\tau} \right)^{rs\tau Kn} \\
&\stackrel{(15)}{=} n \cdot (2^{rs+7}e \cdot Wrs \log(128Wr))^{rs\tau Kn} \\
&\leq n \cdot \exp \left(rs\tau Kn \left(rs + 6 + \ln(Wrs \log(128Wr)) \right) \right) \\
&\stackrel{(15)}{\leq} n \cdot \exp \left(\frac{pn}{128W} \right) \\
&\leq \exp \left(\frac{pn}{64W} \right). \tag{25}
\end{aligned}$$

Finally, combining (23) and (25) with (21) leads to

$$\mathbb{P}([n]_p \in \mathcal{B}) \leq \exp \left(-\frac{pn}{32W} \right) \cdot \exp \left(\frac{pn}{64W} \right) = \exp \left(-\frac{pn}{64W} \right) < \frac{1}{2}.$$

Up to the proof of Fact 4.3 this concludes the proof of Claim 4.2. \square

Proof of Fact 4.3. Recall that $W = \text{vdW}(k; r)$ asserts that $A \rightarrow (AP_k)_r$ for every arithmetic progression $A \subseteq \mathbb{N}$ of length W . Consider an arbitrary $(r+1)$ -colouring of $[n]$. Suppose at most $\frac{n}{4W}$ elements of $[n]$ receive colour $r+1$. From the observation that for every $w \leq W/2$ and every $i \in [n]$ there are at most $\frac{n-1}{W-w}$ distinct AP_W 's in $[n]$ having i at position w or $W-w+1$, one can deduce that every $i \in [n]$ is contained in at most n different AP_W 's. Consequently, there are at least

$$\left| \binom{[n]}{AP_W} \right| - \frac{n^2}{4W} \geq \frac{n^2}{4W}$$

AP_W 's containing no element of the last colour, where we used $\left| \binom{[n]}{AP_W} \right| \geq \frac{n^2}{2W}$ for the last inequality (cf. (18)).

Owing to the choice of W every such r -coloured AP_W contains a monochromatic AP_k in one of the first r colours. On the other hand, every AP_k can be contained in at most $\binom{W}{2}$ different AP_W 's in $[n]$. Therefore, there exist at least

$$\frac{n^2}{4W} \cdot \frac{2}{W^2} = \frac{n^2}{2W^3} \geq \frac{1}{W^3} \cdot \left| \binom{[n]}{AP_k} \right|$$

distinct monochromatic AP_k 's in $[n]$ coloured in one of the first r colours and the fact follows. \square

§5. PROOF OF THEOREM 1.4

In this section we establish that for all integers $r \geq 2$, $g \geq 3$, and $k \geq 3$, there exists a graph H on at most $k^{40gk^4} R^{40gk^2}$ vertices such that $\binom{H}{K_k}$ has chromatic number greater than r and girth at least g , where $R = R(K_k; r)$ is the r -colour Ramsey number for K_k .

Proof. We first define all constants involved in the proof. Given the uniformity $k \geq 3$, the number of colours $r \geq 2$, and the minimum girth $g \geq 3$, we denote by $R = R(K_k; r)$ the r -colour Ramsey number for K_k . In the estimates below we sometimes use the trivial observation $2^r < R$. For a later application of the Container Lemma (Theorem 2.1), we define the involved auxiliary constants (and observe some immediate bounds)

$$\varepsilon = \frac{1}{2r \binom{R}{k}} < \frac{1}{2} \quad \text{and} \quad D_\tau = \left(\frac{6 \cdot \binom{k}{2}! \cdot 2^{\binom{k}{2}} \binom{k}{2} k^k}{\varepsilon} \right)^{10/k^2} < 2^{3k^2/2} k^{20} R^{20/k} \quad (26)$$

and integers

$$K = 800 \binom{k}{2} \left(\binom{k}{2}! \right)^3 < k^{3k^2} \quad \text{and} \quad s = \lceil K \log(1/\varepsilon) \rceil < k^{3k^2} \log(rR^k). \quad (27)$$

We introduce another auxiliary constant

$$D_p = 50R^2 r^2 s^2 K D_\tau \log(50R^2 r) < k^{10k^2+30} R^{5+20/k} \quad (28)$$

and set

$$n = D_p^{k^2(5+g)} < D_p^{3k^2g} < k^{40gk^4} R^{40gk^2}. \quad (29)$$

Finally, we define the following three parameters in terms of some of the constants above

$$\tau = \frac{D_\tau}{n^{2/(k+1)}}, \quad p = \frac{D_p}{n^{2/(k+1)}}, \quad \text{and} \quad t = \frac{p}{2R^2} \binom{n}{2}. \quad (30)$$

Having defined all involved constants we shall show the following two claims, which yield the theorem.

Claim 5.1. *With probability larger than $1/2$, the random graph $G(n, p)$ has the property that there is a set $T \subseteq E(G(n, p))$ of size at most t such that $\text{girth} \left(\binom{G(n, p) - T}{K_k} \right) \geq g$.*

Claim 5.2. *With probability larger than $1/2$, the random graph $G(n, p)$ has the property that $\chi \left(\binom{G(n, p) - T}{K_k} \right) > r$ for every subset $T \subseteq E(G(n, p))$ of size at most t .*

Both claims together show that with positive probability there exists a graph G which contains a set $T \subseteq E(G)$ of size at most t such that $H = G - T$ satisfies $\text{girth} \left(\binom{H}{K_k} \right) \geq g$ and $\chi \left(\binom{H}{K_k} \right) > r$. Consequently, Theorem 1.4 follows from the choice of n in (29). \square

Proof of Claim 5.1. Recall that a 2-cycle is a pair of edges e_1, e_2 such that $|e_1 \cap e_2| > 1$ and for $j > 2$ a j -cycle is a cyclical sequence of j edges e_1, e_2, \dots, e_j where the intersection of two consecutive edges is exactly one i.e. $|e_i \cap e_{i+1}| = 1$ (addition mod j), the intersection of any two nonconsecutive edges is empty, and the intersection points for each pair of consecutive edges is unique.

Define X_j to be the number of j -cycles in the system of copies of K_k in $G(n, p)$. We first work to bound X_2 . If $k = 3$, we trivially have $\mathbb{E}[X_2] = 0$. Otherwise for $k \geq 4$, a 2-cycle corresponds to two copies of K_k that intersect in more than two edges, and thus in more than two vertices. Furthermore, we see that two copies of K_k that intersect in i vertices together span exactly $2k - i$ vertices and $2\binom{k}{2} - \binom{i}{2}$ edges. With this in mind, the following bounds $\mathbb{E}[X_2] < t/4$ in $\binom{G(n,p)}{K_k}$:

$$\begin{aligned} \frac{\mathbb{E}[X_2]}{t/4} &= \frac{8R^2}{p\binom{n}{2}} \cdot \mathbb{E}[X_2] \leq \frac{32R^2}{pn^2} \cdot \sum_{i=3}^{k-1} n^{2k-i} p^{2\binom{k}{2}-\binom{i}{2}} \\ &= 32R^2 n^{2k-2} p^{2\binom{k}{2}-1} \sum_{i=3}^{k-1} n^{(i^2-2i-ki)/(k+1)} D_p^{-\binom{i}{2}} \\ &\leq 32R^2 n^{2k-2} p^{2\binom{k}{2}-1} \cdot k \cdot \max_{3 \leq i \leq k-1} \left\{ n^{(i^2-2i-ki)/(k+1)} \right\} \\ &\leq 32R^2 n^{2k-2} p^{2\binom{k}{2}-1} \cdot k \cdot n^{(3-3k)/(k+1)} \\ &= \frac{32kR^2 D_p^{k^2-k-1}}{n^{(k-3)/(k+1)}} \leq \frac{D_p^{k^2}}{n^{1/5}} \stackrel{(29)}{<} 1. \end{aligned}$$

We now bound $\sum_{j=3}^{g-1} X_j$. For $j > 2$, a j -cycle in $\binom{K_n}{K_k}$ consists of a cyclically ordered set of j copies of K_k such that each two consecutive copies intersect in exactly one edge of K_n . Thus, a j -cycle corresponds to a set of K_k 's in K_n that span at most $kj - 2j$ vertices in K_n and exactly $\binom{k}{2}j - j$ edges in K_n . From this, we see that, for $2 < j < g$, we have

$$\mathbb{E}[X_j] \leq n^{kj-2j} p^{\binom{k}{2}j-j} = \left(n^{k-2} p^{\binom{k}{2}-1} \right)^j = D_p^{(\binom{k}{2}-1)j}.$$

Using this, we establish $\sum_{j=3}^{g-1} \mathbb{E}[X_j] < t/4$:

$$\frac{\sum_{j=3}^{g-1} \mathbb{E}[X_j]}{t/4} \leq \frac{8R^2}{p\binom{n}{2}} \cdot g \cdot D_p^{((\binom{k}{2}-1)g)} \leq \frac{32R^2 g}{pn^2} D_p^{((\binom{k}{2}-1)g)} \leq \frac{D_p^{k^2 g}}{n} < 1.$$

Thus, we have shown $\sum_{j=2}^{g-1} \mathbb{E}[X_j] < t/4 + t/4 = t/2$. By Markov's inequality, this gives that, with probability bigger 1/2, the hypergraph $\binom{G(n,p)}{K_k}$ contains less than t cycles of length less than g . For each such cycle, removing one vertex (which is an edge in $G(n, p)$) concludes the proof of Claim 5.1. \square

Proof of Claim 5.2. The proof relies on an application of the Container Lemma (Theorem 2.1) to the $\binom{k}{2}$ -uniform hypergraph $\mathcal{H} = (K_n, \binom{K_n}{K_k})$. In view of that we will first verify condition (1) for our choices of ε and τ in (26) and (30). Recalling the definition of the average degrees d_j for $j = 1, \dots, \binom{k}{2}$ of \mathcal{H} , we note that

$$d_1 = \binom{n-2}{k-2} \geq \frac{n^{k-2}}{k^k}.$$

For $j \geq 2$, letting k_j be the smallest integer such that $j \leq \binom{k_j}{2}$, we have

$$d_j \leq \binom{n-k_j}{k-k_j} \leq n^{k-k_j}.$$

Consequently, for every $j = 2, \dots, \binom{k}{2}$ this gives

$$\frac{d_j}{d_1 \cdot \tau^{j-1}} \leq \frac{k^k \cdot n^{2-k_j} \cdot n^{\frac{2j-2}{k+1}}}{D_\tau^{j-1}} \leq \frac{k^k \cdot n^{2-k_j} \cdot n^{\frac{2\binom{k_j}{2}-2}{k+1}}}{D_\tau^{j-1}} = \frac{k^k \cdot n^{\frac{(k_j-2)(k_j-k)}{k+1}}}{D_\tau^{j-1}}. \quad (31)$$

For $k_j = k$, i.e., for $j = \binom{k-1}{2} + 1, \dots, \binom{k}{2}$ we, therefore, get

$$\frac{d_j}{d_1 \cdot \tau^{j-1}} \leq \frac{k^k}{D_\tau^{j-1}} \leq \frac{k^k}{D_\tau^{\binom{k-1}{2}}} \leq \frac{k^k}{D_\tau^{k^2/10}}, \quad (32)$$

where we used $D_\tau \geq 1$ and $k \geq 3$ for the last inequalities. For integers $3 \leq k_j \leq k-1$ we note that $k \geq 4$ and $(k_j-2)(k_j-k)$ is maximized for $k_j = 3$ (and $k_j = k-1$). Hence, in this case we can bound the R-H-S in (31) to give

$$\frac{d_j}{d_1 \cdot \tau^{j-1}} \leq \frac{k^k}{n^{1/5}}.$$

Summarizing, since $k_j \geq 3$ for $j \geq 2$ we arrive for $h = \binom{k}{2}$ at

$$\frac{6 \cdot h! \cdot 2^{\binom{h}{2}}}{d_1} \sum_{j=2}^h \frac{d_j}{2^{\binom{j-1}{2}} \tau^{j-1}} \leq \frac{6 \cdot h! \cdot 2^{\binom{h}{2}} \cdot h \cdot k^k}{\min\{D_\tau^{k^2/10}, n^{1/5}\}} \stackrel{(26),(29)}{\leq} \varepsilon$$

and this shows that condition (1) of Theorem 2.1 holds. Consequently, for every independent set $I \subseteq V(\mathcal{H})$ we can apply conclusions (i)–(iii) of the Container Lemma with the constants defined above.

We consider the family \mathcal{B} of all graphs $B \subseteq K_n$ such that there exists a set $T \subseteq E(B)$ of size at most t and $(B-T) \rightarrow (K_k)_r$, i.e., there exists an r -colouring of the edges of the graph $B-T$ without a monochromatic copy of K_k . In other words, $\chi(K_k^{B-T}) \leq r$ and we may view \mathcal{B} as the set of all ('bad') graphs on n vertices that do not have the desired property of Claim 5.2. Below we establish Claim 5.2 by showing

$$\mathbb{P}(G(n, p) \in \mathcal{B}) < \frac{1}{2}.$$

Consider any graph $B \in \mathcal{B}$. By the definition of \mathcal{B} , there exists a set $T^B \subseteq E(B)$ of size $|T^B| \leq t$ and a partition $E(B) \setminus T^B = I_1^B \cup \dots \cup I_r^B$ with the property that none of the sets I_i^B contains a K_k . In particular, each I_i^B is an independent set in \mathcal{H} and, therefore, properties (i)–(iii) of the Container Lemma assert that for every $i \in [r]$ there exists an s -tuple $\mathcal{S}_i^B = (\mathcal{S}_{i,1}^B, \dots, \mathcal{S}_{i,s}^B)$ of subsets of I_i^B and a container set $\mathcal{C}(\mathcal{S}_i^B) \supseteq I_i^B$ such that

$$|\mathcal{S}_{i,\sigma}^B| \leq \tau K \binom{n}{2} \quad (33)$$

for every $\sigma \in [s]$ and

$$|e_{\mathcal{H}}(\mathcal{C}(\mathcal{S}_i^B))| \leq \varepsilon \binom{n}{k}. \quad (34)$$

We also set $\mathcal{S}^B = (\mathcal{S}_1^B, \dots, \mathcal{S}_r^B)$ and $\mathcal{C}^B = (\mathcal{C}(\mathcal{S}_1^B), \dots, \mathcal{C}(\mathcal{S}_r^B))$.

Moreover, for any possible r -tuple $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ of s -tuples of sets of size at most $\tau K \binom{n}{2}$ we consider the corresponding container vector $\mathcal{C}(\mathcal{S}) = (\mathcal{C}(\mathcal{S}_1), \dots, \mathcal{C}(\mathcal{S}_r))$ given by the Container Lemma. We denote by $D(\mathcal{S})$ its complement in $E(K_n)$ given by

$$D(\mathcal{S}) = E(K_n) \setminus (\mathcal{C}(\mathcal{S}_1) \cup \dots \cup \mathcal{C}(\mathcal{S}_r)).$$

We observe that for any $B \in \mathcal{B}$ the following two properties hold:

- (a) $\bigcup_{i \in [r]} \bigcup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}^B \subseteq (E(B) \setminus D(\mathcal{S}^B))$ and
- (b) $|E(B) \cap D(\mathcal{S}^B)| \leq |T^B| \leq t$, since

$$E(B) \setminus T^B = I_1^B \cup \dots \cup I_r^B \subseteq \mathcal{C}(\mathcal{S}_1^B) \cup \dots \cup \mathcal{C}(\mathcal{S}_r^B) = E(K_n) \setminus D(\mathcal{S}^B).$$

From (a) and (b) we infer that

$$\begin{aligned} \mathbb{P}(G(n, p) \in \mathcal{B}) &\leq \sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\bigcup_{i \in [r]} \bigcup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} \cdot \mathbb{P}\left(|E(G(n, p)) \cap D(\mathcal{S})| \leq t\right) \\ &\leq \max_{\mathcal{S}} \mathbb{P}\left(|E(G(n, p)) \cap D(\mathcal{S})| \leq t\right) \sum_{\mathcal{S}} p^{|\bigcup_{i \in [r]} \bigcup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|}, \end{aligned} \quad (35)$$

where the sum and the maximum are taken over all r -tuples $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ of s -tuples $\mathcal{S}_i = (\mathcal{S}_{i,1}, \dots, \mathcal{S}_{i,s})$ of sets $\mathcal{S}_{i,\sigma}$ of size at most $\tau K \binom{n}{2}$ for $i \in [r]$ and $\sigma \in [s]$. We will use property (ii) of the Container Lemma to bound the maximum probability in (35), while our choice of constants allow us to derive a sufficient bound for the sum in (35).

For the maximum probability we first observe that for every $\mathcal{S} = (\mathcal{S}_1, \dots, \mathcal{S}_r)$ we have

$$|D(\mathcal{S})| > \frac{1}{R^2} \binom{n}{2}. \quad (36)$$

For the proof we use the fact that for any $(r+1)$ -colouring of $E(K_n)$ either there are more than

$$\frac{1}{2^{\binom{R}{k}}} \binom{n}{k}$$

monochromatic copies of K_k in the first r colours or there are more than $\frac{1}{R^2} \binom{n}{2}$ edges having the last colour (see, e.g., [25, Proposition 8]).

In view of this fact, we consider $\mathcal{C}(\mathcal{S}_1) \cup \dots \cup \mathcal{C}(\mathcal{S}_r) \cup D(\mathcal{S})$ as an $(r+1)$ -colouring of $E(K_n)$. Owing to property (ii) of the Container Lemma (see (34)), every $\mathcal{C}(\mathcal{S}_i)$ contains at most $\varepsilon \binom{n}{k}$ copies of K_k and, hence, there are at most

$$r \cdot \varepsilon \binom{n}{k} \stackrel{(26)}{=} \frac{1}{2} \binom{n}{k}^{(R)}$$

monochromatic copies in the first r colours. Therefore, the mentioned fact above yields (36). In particular, the choice of t combined with (36) yields $t < p|D(\mathcal{S})|/2$ and, consequently, Chernoff's inequality (see, e.g., [16, Theorem 2.1]) asserts

$$\mathbb{P}\left(|E(G(n, p)) \cap D(\mathcal{S})| \leq t\right) \leq \exp\left(-\frac{p}{8R^2} \binom{n}{2}\right) \quad (37)$$

for every \mathcal{S} considered here. In particular, (37) bounds the maximum probability considered in the R-H-S of (35) and below we turn to the sum in (35).

Owing to $|\mathcal{S}_{i,\sigma}| \leq \tau K \binom{n}{2}$ for every $i \in [r]$ and $\sigma \in [s]$ we have

$$\sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} \leq \sum_{m=0}^{r \cdot s \cdot \tau K \binom{n}{2}} \binom{\binom{n}{2}}{m} 2^{rsm} p^m \leq \sum_{m=0}^{r \cdot s \cdot \tau K \binom{n}{2}} \left(\frac{e \binom{n}{2}}{m} 2^{rs} p\right)^m.$$

Since the function $m \mapsto (e \binom{n}{2} 2^{rs} p/m)^m$ is unimodal and attains its maximum value for $m_0 = \binom{n}{2} 2^{rs} p \geq r s \tau K \binom{n}{2}$ (see (28)), we can bound the summands in R-H-S above by the last one and obtain

$$\begin{aligned} \sum_{\mathcal{S}=(\mathcal{S}_1, \dots, \mathcal{S}_r)} p^{|\cup_{i \in [r]} \cup_{\sigma \in [s]} \mathcal{S}_{i,\sigma}|} &\leq \left(r s \tau K \binom{n}{2} + 1\right) \cdot \left(\frac{e \binom{n}{2}}{r s \tau K \binom{n}{2}} 2^{rs} p\right)^{r s \tau K \binom{n}{2}} \\ &\leq n^2 \cdot \left(\frac{2^{rs} e D_p}{r s K D_\tau}\right)^{r s \tau K \binom{n}{2}} \\ &\stackrel{(28)}{=} n^2 \cdot (2^{rs} e \cdot 50R^2 r s \log(50R^2 r))^{r s \tau K \binom{n}{2}} \\ &\leq n^2 \cdot \exp\left(r s \tau K \binom{n}{2} (r s + 1 + \ln(50R^2 r s \log(50R^2 r)))\right) \\ &\stackrel{(28)}{\leq} n^2 \cdot \exp\left(\frac{p}{16R^2} \binom{n}{2}\right) \\ &\leq \exp\left(\frac{p}{12R^2} \binom{n}{2}\right). \end{aligned} \quad (38)$$

Finally, combining (37) and (38) with (35) leads to

$$\mathbb{P}(G(n, p) \in \mathcal{B}) \leq \exp\left(-\frac{p}{8R^2} \binom{n}{2}\right) \cdot \exp\left(\frac{p}{12R^2} \binom{n}{2}\right) = \exp\left(-\frac{p}{24R^2} \binom{n}{2}\right) < \frac{1}{2}$$

and Claim 5.2 follows. \square

§6. RAMSEY GRAPHS USING EXTREMAL CONSTRUCTIONS

Recall that in Theorem 1.2 we showed for all integers k and $r \geq 2$ that

$$f(2k, r) = \min \{|V(H)| : \text{girth}(H) = 2k \text{ and } H \longrightarrow (C_{2k})_r\} \leq \exp(c_2 k^2 (\log r + k \log k)).$$

For three small cases of k , we are able to deduce better bounds for $f(2k, r)$ using well known extremal constructions of graphs with girth 6, 8, and 12, respectively.

Theorem 6.1. *We have $f(6, r) = O(r^6)$, $f(8, r) = O(r^{12})$, and $f(12, r) = O(r^{30})$.*

Before proving Theorem 6.1, we first introduce some notation and state an observation upon which the proof is based. Let $\text{ex}(n; C_k)$ denote the maximum number of edges in an n vertex graph that does not contain a cycle of length k . Similarly, let $\text{ex}(n; C_3, C_4, \dots, C_{k-1})$ denote the maximum number of edges in a graph with girth k .

Fact 6.2. *If $\text{ex}(n; C_3, C_4, \dots, C_{2k-1}) > r \cdot \text{ex}(n; C_3, C_4, \dots, C_{2k})$, then $f(2k, r) \leq n$.*

Indeed, by definition of the extremal function there exists a graph G on n vertices with girth $2k$ that has $\text{ex}(n; C_3, C_4, \dots, C_{2k-1})$ edges. Clearly, every r -colouring of G yields a monochromatic subgraph with at least $\text{ex}(n; C_3, C_4, \dots, C_{2k-1})/r > \text{ex}(n; C_3, C_4, \dots, C_{2k})$ edges, which must contain a monochromatic C_{2k} since the monochromatic subgraph still has girth at least $2k$.

Proof of Theorem 6.1. To make use of this fact to prove Theorem 6.1, we use the result of Erdős and Simonovits from [13] that for every positive integer k , we have

$$\text{ex}(n; C_3, C_4, \dots, C_{2k+1}) = O(n^{1+1/k}).$$

Since any graph contains a bipartite subgraph with half of its edges we have

$$\begin{aligned} \text{ex}(n; C_3, C_4, C_5, C_6, \dots, C_{2k}) &\leq \text{ex}(n; C_4, C_6, \dots, C_{2k}) \\ &\leq 2 \cdot \text{ex}(n; C_3, C_4, C_5, C_6, \dots, C_{2k+1}) = O(n^{1+1/k}). \end{aligned} \quad (39)$$

Erdős and Simonovits conjectured in [13] that for every positive integer $k \geq 2$,

$$\text{ex}(n; C_3, C_4, \dots, C_{2k-1}) = \Omega(n^{1+1/(k-1)}). \quad (40)$$

This has been observed for $k = 3$ by Klein (see [6]) and follows for $k = 4$ by the work of Singleton [28], and for $k = 6$ by the work of Benson [2]. For $k \in \{3, 4, 6\}$, inequalities (39) and (40) give that

$$\text{ex}(n; C_3, C_4, \dots, C_{2k-1}) = \Omega(n^{1+1/(k-1)}) > r \cdot O(n^{1+1/k}) = r \cdot \text{ex}(n; C_3, C_4, \dots, C_{2k}),$$

holds, provided that

$$n \geq \tilde{c} r^{k(k-1)},$$

for some sufficiently large constant \tilde{c} . Consequently, Fact 6.2 yields

$$f(2k, r) \leq n = \Omega(r^{k(k-1)})$$

for $k \in \{3, 4, 6\}$ and the theorem follows. \square

We remark that establishing (40) for all k , implies $f(2k, r) = O(r^{k(k-1)})$ for all k by the same argument.

REFERENCES

- [1] J. Balogh, R. Morris, and W. Samotij, *Independent sets in hypergraphs*, J. Amer. Math. Soc. **28** (2015), no. 3, 669–709, DOI [10.1090/S0894-0347-2014-00816-X](https://doi.org/10.1090/S0894-0347-2014-00816-X). MR3327533 [↑1.4, 2](#)
- [2] C. T. Benson, *Minimal regular graphs of girths eight and twelve*, Canad. J. Math. **18** (1966), 1091–1094. MR0197342 (33 #5507) [↑6](#)
- [3] T. F. Bloom, *A quantitative improvement for Roth’s theorem on arithmetic progressions*, J. Lond. Math. Soc. (2) **93** (2016), no. 3, 643–663. MR3509957 [↑1.2](#)
- [4] J. A. Bondy and P. Erdős, *Ramsey numbers for cycles in graphs*, J. Combinatorial Theory Ser. B **14** (1973), 46–54. MR0317991 (47 #6540) [↑1.1](#)
- [5] J. A. Bondy and M. Simonovits, *Cycles of even length in graphs*, J. Combinatorial Theory Ser. B **16** (1974), 97–105. MR0340095 (49 #4851) [↑1.1](#)
- [6] P. Erdős, *On sequences of integers no one of which divides the product of two others and on some related problems*, Mitt. Forsch.-Inst. Math. und Mech. Univ. Tomsk **2** (1938), 74–82. [↑6](#)
- [7] ———, *Extremal problems in graph theory*, Theory of Graphs and its Applications (Proc. Sympos. Smolenice, 1963), Publ. House Czechoslovak Acad. Sci., Prague, 1964, pp. 29–36. MR0180500 (31 #4735) [↑1.1](#)
- [8] P. Erdős, *Problems and results on finite and infinite graphs*, Recent advances in graph theory (Proc. Second Czechoslovak Sympos., Prague, 1974), Academia, Prague, 1975, pp. 183–192. MR0389669 (52 #10500) [↑1.1, 1.3](#)
- [9] ———, *Problems and results in combinatorial number theory*, Journées Arithmétiques de Bordeaux (Conf., Univ. Bordeaux, Bordeaux, 1974), Soc. Math. France, Paris, 1975, pp. 295–310. Astérisque, Nos. 24–25. MR0374075 (51 #10275) [↑1.2](#)
- [10] P. Erdős and R. L. Graham, *On partition theorems for finite graphs*, Infinite and finite sets (Colloq., Keszthely, 1973; dedicated to P. Erdős on his 60th birthday), Vol. I, North-Holland, Amsterdam, 1975, pp. 515–527. Colloq. Math. Soc. János Bolyai, Vol. 10. MR0373959 (51 #10159) [↑1.1](#)
- [11] P. Erdős and A. Hajnal, *Reserach Problems*, Acta Math. Acad. Sci. Hungar **17** (1966), 61–99. MR0193025 (33 #1247) [↑1.2](#)
- [12] ———, *Research problems 2-5*, J. Combinatorial Theory **2** (1967), 104–105. [↑1.3](#)
- [13] P. Erdős and M. Simonovits, *Compactness results in extremal graph theory*, Combinatorica **2** (1982), no. 3, 275–288, DOI [10.1007/BF02579234](https://doi.org/10.1007/BF02579234). MR698653 (84g:05083) [↑6, 6](#)

- [14] J. Folkman, *Graphs with monochromatic complete subgraphs in every edge coloring.*, SIAM J. Appl. Math. **18** (1970), 19–24. MR0268080 (42 #2979) ↑1.3
- [15] R. L. Graham and J. Nešetřil, *Large minimal sets which force long arithmetic progressions*, J. Combin. Theory Ser. A **42** (1986), no. 2, 270–276, DOI 10.1016/0097-3165(86)90097-X. MR847557 (88c:11015) ↑1.2
- [16] S. Janson, T. Łuczak, and A. Ruciński, *Random graphs*, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2000. MR1782847 (2001k:05180) ↑3, 4, 5
- [17] M. Jenssen and J. Skokan, *Exact Ramsey numbers of odd cycles via nonlinear optimisation*. <https://arxiv.org/abs/1608.05705>. ↑1.1
- [18] R. Nenadov and A. Steger, *A short proof of the random Ramsey theorem*, Combin. Probab. Comput. **25** (2016), no. 1, 130–144, DOI 10.1017/S0963548314000832. ↑2, 3
- [19] J. Nešetřil and V. Rödl, *The Ramsey property for graphs with forbidden complete subgraphs*, J. Combinatorial Theory Ser. B **20** (1976), no. 3, 243–249. MR0412004 (54 #133) ↑1.3
- [20] ———, *Van der Waerden theorem for sequences of integers not containing an arithmetic progression of k terms*, Comment. Math. Univ. Carolinae **17** (1976), no. 4, 675–681. MR0441906 (56 #297) ↑1.2
- [21] ———, *Simple proof of the existence of restricted Ramsey graphs by means of a partite construction*, Combinatorica **1** (1981), no. 2, 199–202, DOI 10.1007/BF02579274. MR625551 (83a:05101) ↑1.3
- [22] ———, *Sparse Ramsey graphs*, Combinatorica **4** (1984), no. 1, 71–78, DOI 10.1007/BF02579159. MR739415 (85h:05073) ↑1.3
- [23] V. Rödl, *On Ramsey families of sets*, Graphs Combin. **6** (1990), no. 2, 187–195, DOI 10.1007/BF01787730. MR1073689 (91m:05137) ↑1.2
- [24] V. Rödl and A. Ruciński, *Threshold functions for Ramsey properties*, J. Amer. Math. Soc. **8** (1995), no. 4, 917–942, DOI 10.2307/2152833. MR1276825 (96h:05141) ↑1.1, 1.2
- [25] V. Rödl, A. Ruciński, and M. Schacht, *An exponential-type upper bound for Folkman numbers*, Combinatorica **37** (2017), no. 4, 767–784. MR3694711 ↑2, 3, 3, 5
- [26] T. Sanders, *On Roth’s theorem on progressions*, Ann. of Math. (2) **174** (2011), no. 1, 619–636, DOI 10.4007/annals.2011.174.1.20. MR2811612 (2012f:11019) ↑1.2
- [27] D. Saxton and A. Thomason, *Hypergraph containers*, Invent. Math. **201** (2015), no. 3, 925–992, DOI 10.1007/s00222-014-0562-8. MR3385638 ↑2, 2
- [28] R. Singleton, *On minimal graphs of maximum even girth*, J. Combinatorial Theory **1** (1966), 306–332. MR0201347 (34 #1231) ↑6
- [29] J. Spencer, *Restricted Ramsey configurations*, J. Combinatorial Theory Ser. A **19** (1975), no. 3, 278–286. MR0382058 (52 #2946) ↑1.2

DEPARTAMENTO DE MATEMÁTICA Y CIENCIA DE LA COMPUTACIÓN, UNIVERSIDAD DE SANTIAGO DE CHILE, CHILE

Email address: `hiep.han@usach.cl`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, USA

Email address: `{tretter | rodl}@mathcs.emory.edu`

FACHBEREICH MATHEMATIK, UNIVERSITÄT HAMBURG, HAMBURG, GERMANY

Email address: `schacht@math.uni-hamburg.de`